ORAL BIOLOGY AND MEDICAL TECHNOLOGY

ISBN:
978-625-90649-0-1

ORAL BIOLOGY AND MEDICAL TECHNOLOGY-2026

ISBN: 978-625-90649-0-1

DOI: 10.5281/zenodo.21194891

July / 2026

Ankara / Türkiye

Copyright © 2026 by ISPEC publishing house

All rights reserved. No part of this publication may be reproduced, distributed or transmitted in any form or by any means, including photocopying, recording or other electronic or mechanical methods, without the prior written permission of the publisher, except in the case of brief quotations embodied in critical reviews and certain other noncommercial uses permitted by copyright law. UBAK International Academy of Sciences Association Publishing House®

(The Licence Number of Publicator: 2014/31220)

E mail: info@ispecbooks.com

www.ispecbooks.com

It is responsibility of the author to abide by the publishing ethics rules.

ISPEC Publishing House – 2026©

ISBN: 978-625-90649-0-1

July / 2026

Ankara / Türkiye

ORAL BIOLOGY AND MEDICAL TECHNOLOGY

AUTHORS
Prof. Dr. Mehwash KASHIF
Prof. Dr.Vıjaya LAXMI
Assoc. Prof. Dr. Muhammad Abdullah KAMRAN
Dr. Maqubool HOSAIN
Dr. Naveen TRIVEDI
Dr. Kainat ARSHAD
Dr. Erum FATIMA
Mila POPOVIĆ
Svetozar ĐURANOVIĆ
Marko POPOVIĆ

TABLE OF CONTENTS

PREFACE.....i

CHAPTER 1
MULTIDISCIPLINARY OBSERVATION LIKE BIOLOGICAL INFLUENCES, AN EXAMPLE OF PROBIOTICS IN THE MANAGEMENT OF THE TREATMENT OF DENTAL PATIENTS
Mila POPOVIĆ
Svetozar ĐURANOVIĆ
Marko POPOVIĆ.....1

CHAPTER 2
FUNDAMENTALS OF ORAL CAVITY BIOLOGY
Prof. Dr. Mehwash KASHIF
Assoc. Prof. Dr. Muhammad Abdullah KAMRAN
Dr. Kainat ARSHAD
Dr. Erum FATIMA 17

CHAPTER 3
INTERNET OF THINGS (IOT) IN MEDICAL SCIENCES: APPLICATIONS, CHALLENGES, AND FUTURE PERSPECTIVES
Dr. Maqubool HOSAIN
Dr. Naveen TRIVEDI
Prof. Dr.Vıjaya LAXMI. 55

PREFACE

Oral health is closely connected to general health, biological processes, preventive care, and technological development. Contemporary research increasingly examines the role of oral biology, microbiological balance, and digital health technologies in improving diagnosis, treatment, and patient care. These developments highlight the importance of interdisciplinary approaches that connect dental sciences with broader advances in medical research.

The chapters in this volume examine selected topics at the intersection of oral health and medical technology. The first chapter discusses the potential role of probiotics in the management and treatment of dental patients, highlighting the relevance of biological influences in oral care. The second chapter presents the fundamental principles of oral cavity biology and its importance for understanding oral health and disease. The third chapter explores the applications, challenges, and future perspectives of the Internet of Things in medical sciences.

By bringing together studies on oral biology, probiotic approaches, and digital medical technologies, this volume reflects the evolving nature of healthcare research. The chapters demonstrate how biological knowledge and technology-supported systems can contribute to more effective, preventive, and patient-centered health services.

It is hoped that this book will serve as a useful academic resource for researchers, students, healthcare professionals, and scholars interested in dentistry, oral biology, medical sciences, digital health, and emerging healthcare technologies. We extend our sincere appreciation to all contributing authors for their valuable scholarly efforts.

Editorial Team
July 2026, Türkiye

CHAPTER 1
MULTIDISCIPLINARY OBSERVATION LIKE BIOLOGICAL INFLUENCES, AN EXAMPLE OF PROBIOTICS IN THE MANAGEMENT OF THE TREATMENT OF DENTAL PATIENTS

[1]Mila POPOVIĆ
[2]Svetozar ĐURANOVIĆ
[3]Marko POPOVIĆ

[1]Dentist, Medical Faculty of Novi Sad, Serbia, slobodan.popovic49@gmail.com, ORCID ID: 0009-0007-8426-0246
[2]Doctor of Medicine, Clinical Center of Vojvodina, Novi Sad, Serbia, gandra.dj@hotmail.com, ORCID ID: 0009-0004-6221-1353
[3]University of Novi Sad, Serbia, vera.popovic@ifvcns.ns.ac.rs, ORCID ID: 0009-0000-2149-9750

INTRODUCTION

Multidisciplinary approach, which includes numerous disciplines such as applied biology, patient care, and dental services, should be a complete observation that, within the framework of medical services (Dragosavac et al., 2023; Bakmaz et al., 2024-a), interconnects and strengthens numerous segments of action, from legal regulation (Jovanović et al., 2025; Lutovac et al., 2025; Bakmaz et al., 2025) to the application of biologically successful treatment in the field of dentistry (Popović, 2025).

Application in numerous treatments requires that applied biology, patient treatment and therefore also within dental services (Newman et al., 2012; Vihnanek, 2015; Dessuane et al., 2018; Dimitrijević, 2019) be connected with real business functions with the aim of achieving a business that has very strong and focused results on action with the aim of improving future economic activities that, in such an observation, relate to dental treatment, primarily economically profitable (Popović et al., 2017; Radović et al., 2019; Vitomir et al., 2020; Arnautović et al., 2022; Tomas-Miskin et al., 2022; Bakmaz et al., 2024) both from the point of view of treatment and from the point of view of patient satisfaction.

One of the important parts of biologically predictable action in the treatment of patients refers to the observation of the conditions created by the state in terms of the action and manifestation of the action of society on the health system, which achieves the basis of multidisciplinary, for example, observed through the action of biological influence, treatment, application of demonstrable substances in treatments that can improve treatment, for example the application of probiotics (Newman and Carranza, 2006) and similar (Tošković et al., 2016; Popović et al., 2021: Radović et al., 2923).

The effect of multidisciplinary approach, which includes numerous disciplines such as applied biology, patient treatment, dental services, is a great, but not exclusive effect, because the totality of the impact includes satisfying the connection of biological influences on treatment in, for example, dentistry (Stanišić, 2021) with the socially observed health system (Vitomir et al., 2021; Radović et al., 2021-a), and such observation requires the best and most optimal operation of both the biological-medical profession and the connection with the entire health system in the country.

Multidisciplinary observation of the action of biology, treatment and return of patients, for example after dental services (Đajić et al., 2001) requires heterogeneous direct action and the creation of numerous impacts on the entire society, which is indisputable. Such action should be comprehensive and continuous, with the focus primarily on creating a lasting connection between biological medical action on the speed of recovery and the achieved effect of treatment in quality that meets the achieved world standards in the future, so that decision-makers can make valid management decisions (Arnautović et al., 2022-a; Vitomir et al., 2023; Bakmaz et al., 2024-b; Popović et al., 2025), primarily within the state's healthcare system.

1. CREATION OF GENERAL OPERATING CONDITIONS FOR THE FUNCTIONING OF A MULTIDISCIPLINARY FORM OF ACTION THROUGH BIOLOGICAL-MEDICAL INFLUENCE ON THE TREATMENT OF PATIENTS

According to the World Health Organization, the creation of general operating conditions for the functioning of a multidisciplinary form of action through biological-medical influence on the treatment of patients is very widely represented and diverse.

One of the noticeable factors of influence on the health care system is the observation of the system of action of the biological-medical influence on the treatment of patients through the created system on the oral health of the population.

Oral observation of the impact of the population represents the state of the mouth, teeth and orofacial structures that enable the individual to perform essential functions, such as breathing, speaking, eating, and contribute to the preservation of self-confidence and the ability to socialize and function without pain.

Only healthy patients, i.e. a population where there is a built health system in which oral health varies during different life ages and is closely related to general health, where the system is seen as a health system that will return the patient to a normal life in the shortest possible period.

Periodontitis is one of the most common chronic diseases of the oral cavity and is the main cause of tooth loss in adults. In such conditions of existence, in a short period of time, the citizens, that is, the population, should be returned to a normal and satisfied life.

Its etiology is complex and involves the interaction of microbiological, immunological and host factors, whereby bacterial infection plays a key role in the initiation and progression of the disease. Biological treatments can mean speeding up the treatment and achieving quick satisfaction of the patient or the population.

The traditional approach to the treatment of periodontitis is based on the mechanical removal of dental plaque and calculus, with the eventual application of antimicrobial agents.

However, the application of probiotics, which by their action on the oral microflora can contribute to the establishment of microbiological balance and reduce the inflammatory response, is gaining increasing importance in modern therapy.

Research in the last decade points to the potential of probiotics as an auxiliary agent in the therapy of periodontitis, which opens the possibility for the development of new, biologically acceptable approaches in the treatment and prevention of this disease.

2. ANATOMY OF PERIODONTICIUM IN THE FOCUS OF OBSERVATION OF BIOLOGICAL MEDICAL TREATMENT

The periodontium is the supporting apparatus of the teeth, and its main function is the fixation of the teeth in the bone amortization of chewing forces.

It consists of four basic components:

- Gingiva ~ mucosa that covers the alveolar bone and surrounds the neck of the tooth. It is divided into: free, attached and interdental gingiva.
- Periodontal ligament A thin connective tissue structure located between the cementum of the root of the tooth and the alveolar bone.
- Tooth cement A thin, mineralized layer that covers the root of a tooth. Fibers are attached to it periodontal ligament.

- Alveolar bone The part of the maxilla and mandible that forms the dental cup (alveolus) in which the tooth is located.

Depending on the localization of the pathological process in the periodontium, two should be distinguished forms of disease.

Gingivitis is a disease caused by the occurrence and development of inflammation exclusively in gingiva, with the presence of dental plaque as the main etiological factor.

Clinically, it is the gingiva red in color, it is enlarged due to edema or fibrosis, the amount of exudate is increased and it bleeds provocation and/or spontaneously.

Inflammation can be of different types and variable intensity.

If the etiological factor of the accompanying changes is detected and successfully removed in time pathological conditions in the gingiva are reversible.

However, in case of non-undertaking of therapeutic measure, gingivitis can damage deeper periodontal tissues and progress to periodontitis.

The host's immune response, innate and acquired, results the release of inflammatory markers such as cytokines, chemokines and matrix metal proteinases.

This defensive reaction leads to the destruction of the supporting apparatus of the tooth.

Based on this, it is concluded that it plays a key role in the pathogenesis of this disease the interaction of bacterial infection and the response of the host's immune system, on which it depends degree of destruction of supporting tissues.

Without bacteria and their products there is no initiation infection, however, the immune system of the host is responsible for the further course of the pathological process.

3. APPRECIATION OF THE IMPORTANCE OF THE EXISTENCE OF A CLINICAL PICTURE OF PERIODONTOPATHY IN THE OVERALL OBSERVATION OF THE PATIENT-HEALTH SYSTEM RELATIONSHIP

The clinical picture is a set of symptoms and signs that can be observed during the examination and that indicate the existence and stage of the disease.

Regardless of the type of disease, the symptoms are similar, although the clinical manifestations are most pronounced in the stage of the full clinical picture of the disease (if we exclude the terminal stage).

The disease is characterized by the following symptoms:

- Gingival inflammation. It implies a change in color, shape and size, consistency, surface structure and subjective complaints of patients. During periodontitis, inflammation occurs in the gingiva, most often of the exudative type.The earliest signs of inflammation are an increased amount of gingival fluid and bleeding during gentle probing.
- Gingival recession.Gum recession, or gingival recession, is the exposure of the tooth root by apical movement of the gingiva. In addition to inflammation of the gingiva, as the most important factor, aging, anomalies of the gingiva and alveolar bone, mucogingival anomalies, orthodontic anomalies, and poor brushing technique also contribute to receding gums.
- Periodontal pockets. A periodontal pocket is a pathological formation and is a pathognomonic symptom of periodontopathy. It occurs as a result of the deepening of the gingival sulcus.

The formation of a periodontal pocket is conditioned by the destruction of periodontal tissues and the apical migration of the attachment epithelium.

The walls of the pocket consist of inflamed gingiva (soft wall) and the root of the tooth (hard wall), the bottom is always more apical 12 than the cement-enamel border and is formed by degenerated attachment epithelium.

The natural opening of the pocket is located in the region of the edge of the inflamed gingiva. Given that periodontal pockets are a pathognomonic symptom of periodontitis in the differential diagnostic procedure, it is necessary to establish their existence, localization, type, determine their depth and, what is of particular importance, determine the level of the connective epithelium.

The depth of the periodontal pocket is the measured distance from the edge of the gums to the bottom of the periodontal pocket, expressed in millimeters, using a periodontal probe.

It is not a dominantly significant parameter for assessing the condition of periodontal tissues, given that it is determined by the extent of destruction of periodontal tissues, but also by the type and intensity of inflammation in the soft wall of the pocket.

We can have high values for the depth of the pocket (due to the enlargement of the gingiva), while the extent of destruction of the deeper periodontal tissues is small.

Conversely, if destructive processes are expressed in all periodontal tissues, due to the loss of gingival tissue and receding gums, the extent of destruction of deeper periodontal tissues can be large, while the values for the depth of the pockets are small.

The key element in assessing the condition of the periodontal tissues is the level of the attachment epithelium, i.e. measured distance from the cement-enamel border to the coronary end of the attachment epithelium.

In people with a healthy periodontium, as well as in those suffering from gingivitis (gingival pockets), the value of the level of the attachment epithelium is always zero.

In patients with periodontitis in the region of the periodontal pockets, the value of the attachment epithelium is always greater than zero.

4. COMPREHENSIVENESS OF THE OBSERVATION OF PROBIOTICS IN THE FRAMEWORK OF THE BIOLOGICAL-MEDICAL SETTING OF POPULATION TREATMENT MANAGEMENT IN THE HEALTH SYSTEM

The name probiotic comes from the Greek word meaning "for life" (Greek pro bios), and the term probiotic received its clear definition in 2003.

Probiotics are living microorganisms that, when ingested in adequate amounts, bring health benefits to the host.

They have been used in medicine for many years for the treatment of inflammatory bowel diseases, the prevention of allergies, the treatment of vaginal infections and the prevention of respiratory tract infections.

According to the general definition of probiotics, it is clear that a large number microorganism can exhibit probiotic properties. Some of the most important representatives of these microorganisms.

When it comes to nutrition, lactic acid bacteria are of special value, among which the most important in the probiotic sense are species from the genera Lactobacillus and catalase-negative species that produce lactic acid as the main product. although phylogenetically, it has many specific and isolated metabolic pathways.

Prebiotics are non-digestible dietary supplements that selectively stimulate the growth and activity of beneficial bacteria in the body, especially the genera Lactobacillus and Bifid bacterium, and at the same time inhibit the growth and activity of potentially harmful bacteria.

Symbiotic are dietary supplements that combine probiotics and prebiotics, which are considered to work together through synergism. Studies have shown that symbiotic can be more effective than when these supplements are used individually.

Antibiotics are important drugs used to treat bacterial infections. However, they can eliminate not only harmful, but also beneficial bacteria that contribute to maintaining health and can strengthen immunity, thereby improving the body's defense against pathogens.

The elimination of beneficial bacteria often leads to digestive problems, most often diarrhea. Antibiotics are the opposite of probiotics. Probiotics represent live microorganisms, most often bacteria, which, when properly applied, have a beneficial effect on the host's health, while antibiotics act bactericidal or bacteriostatic ally, leading to inhibition growth or elimination of bacteria.

Probiotics have a wide range of effects that differ depending on the probiotic strain. In most cases, the combination of activities of several strains is responsible for the mechanism of action, which is why explaining their action is a very complex task.

The therapeutic effect is based on the establishment of microbiological balance. A probiotic must fulfill several conditions in order to benefit oral and general human health. It is necessary for it to survive the conditions in the gastrointestinal tract and to reach the site of action alive.

They must be able to survive in the presence of bile and gastric juice, adhere to epithelial surfaces, remain present in the human GIT, stimulate the immune system and exhibit antagonistic activity against intestinal pathogens such as Clostridium difficile, Salmonella spp., and Helicobacter pylori, which actually means the production of antibacterial substances, i.e. the production of primary metabolites: lactic acid, acetaldehyde, and hydrogen peroxide.

Probiotics are characterized by the ability to adhere to nutrients, primarily indigestible carbohydrates in the large intestine.

Adhesion can be specific (refers to the binding of adhesion to the surface of bacterial cells on the intestinal epithelium) and non-specific. Non-specific adhesion 29 involves electrostatic and hydrophobic interactions that are often the initial stage that precedes specific adhesion.

Also, probiotics have the ability to modify metabolic processes in the GIT. They can increase the activities of some enzymes (e.g. ß-galactosidase which increases lactose tolerance) and decrease the activities of enzymes involved in carcinogenic processes such is nitro reductase and possibly other forms.

The effect on the host is reflected in the modulation of the innate and acquired immune response, by reducing the production of pro-inflammatory cytokines: TNFα, IL-1β and IL-6 and increasing the production of anti-inflammatory cytokines.

They also participate in the development and regulation of the immune system, the synthesis of amino acids, proteins and various vitamins, the absorption of calcium, iron and vitamin D.

Thanks to the aforementioned properties, probiotics are increasingly being introduced into preventive and therapeutic protocols in dentistry, especially in areas where symbiosis is one of the key components of pathogenesis including periodontopathy.

In the treatment of dental diseases, probiotics have been used for the last ten years.

In addition to periodontal disease, they are also used in the treatment of caries and bad breath. When applied in an adequate amount, they have a beneficial effect on the host's health. Studies have shown that the combination of different types of probiotic strains leads to a synergistic enhancement of their 31 individual effects.

In medicine and dentistry, strains from the genera Lactobacillus, Bifid bacterium, Streptococcus, as well as some yeasts such as Saccharomyces is most often used. In modern dentistry, probiotics are increasingly used in the form of various oral preparations, such as toothpastes, tablets, chewing gums and solutions for rinsing the oral cavity.

It has been shown that pastes enriched with probiotics can significantly reduce the presence of Streptococcus mutants, one of the main bacteria responsible for the development of caries.

Local application of these preparations gives results that are comparable to systemic use. Research indicates that probiotic chewing gums can act on the same microorganism, but also contribute to improving the clinical status of patients with periodontal diseases.

Chewing gums and antiseptic solutions with probiotic cultures have also been shown to be effective in reducing bad breath, primarily by inhibiting the formation of sulfur compounds.

Products with Lactobacillus reutter, whether in the form of chewing gum or tablets, can significantly reduce the concentration of S. mutants in saliva.

Probiotics represent a relatively new area within oral medicine. Although the results so far indicate a significant therapeutic potential, additional clinical trials are needed to enable the development of more effective probiotic formulations for the prevention and treatment of oral cavity diseases.

The first sign of periodontal disease is bleeding during probing, while discoloration, swelling, periodontal pockets and, in more advanced stages, luxation and tooth migration are other symptoms.

Current research indicates that probiotics can play a significant role in lowering the 32 pH value of saliva, the production of antioxidant substances that inhibit the growth of bacteria and prevent the formation of dental plaque, as well as in the synthesis of lactic acid with an antibacterial effect, hydrogen peroxide with an antiseptic effect, and antiviral and antifungal agents that contribute to the suppression of pathogenic microorganisms.

In this way, probiotics can play a role in the therapy of periodontal diseases.

Probiotics that are applied for the purpose of improving oral health must be adapted to their form of application.

Unlike those in tablets or capsules for gastrointestinal administration that are swallowed, probiotics for the oral cavity come in the form of means that allow them to be retained in the mouth.

So, for example, one of the most commonly used probiotics for oral administration on the market is available in the form of lozenges that slowly dissolve in the mouth, while preparations such as probiotic mouthwash solutions or sub gingival application in periodontal pockets are also used for research purposes.

Probiotics can never replace the mechanical removal of plaque, but research has shown their potential positive effect as an addition to classic periodontal therapy.

CONCLUSION

Multidisciplinary in many areas of life and work, but also essential being, which includes numerous disciplines such as applied biology, treatment of patients, dental services should be a complete observation, that is, observation should be within the scope of medical services in the process of using the application of biologically successful treatment, which the authors set in the paper to be reflected in the domain of dentistry.

The application of the biological-medical approach in numerous treatments requires the application of biology, the treatment of patients, and thus also within dental services, which unites the connection with the real functions of being within the health system.

The goal of such business with the goal of achieving the use of biologically successful treatment in working with patients is the quick and safe return of the patient, i.e. the citizen, to normal life processes after treatment.

Multidisciplinary in many areas of life and work, but also essential being within the framework of action on the dental turn of research in focus refers primarily to economically profitable treatment both from the point of view of treatment and from the point of view of overall patient satisfaction after the biological-medical treatment.

All of this can be seen as part of the overall biologically predictable action in the treatment of patients, and the entire process of multidisciplinary that the authors of the study point to refers to the observation of the conditions created by the state regarding the action and manifestation of society's action on the health system.

Such activities are not random, but represent well-designed influences with the aim of achieving the economy of treatment of large parts of the population in the country.

The effect of multidisciplinary, which includes numerous disciplines such as applied biology, patient treatment, and dental services, is a large, but not exclusive effect, which is emphasized in this paper through the impact of the use of probiotics as a biological-medical way of improving treatment and creating better treatment effects in the shortest possible time.

The authors pointed out that the application of probiotics and their reflection is aimed at improving the oral health of the general population, where all processes must be adjusted according to the procedure prescribed by the form of application.

Unlike those in tablets or capsules for gastrointestinal administration that are swallowed, probiotics for the oral cavity come in the form of means that allow them to be retained in the mouth. We emphasize that one of the most commonly used probiotics for oral administration on the market is available in the form of lozenges that slowly dissolve in the mouth, while preparations such as probiotic mouthwash solutions or sub gingival application in periodontal pockets are also used for research purposes.

REFERENCES

Arnautović, I., Davidov, T., Nastić, S. & Popović, S. (2022).The importance of making rational business decisions of top management in agricultural enterprises in the Republic of Serbia, Agricultural machinery, 1-8.

Arnautović, I., Davidov, T., Nastić, S. & Popović, S. (2022-a).The importance of business decision-making by top management in agricultural enterprises that use the practical application of new approaches to software solutions in the IT sector in the Republic of Serbia, Agricultural machinery, 51-57.

Bakmaz, O., Dragosavac, M., Brakus, A., Krstajić, G., Borojević, N., Grublješić, Ž., Popović, D. & Popović, S. (2024).The importance of safe financial reporting with the aim of realistic management of agricultural machinery, - example of the Republic of Serbia, Agricultural machinery, 1-6.

Bakmaz, O., Dragosavac, M., Brakus, A., Radaković, M., Arnautović, I., Samardžić, V., Krstajić, G. & Popović, S. (2024.a). Financial security and invoicing in management of public enterprises whose founders are local self-government units, example Republic of Serbia, Lex localis-Journal of Local Self-Government, 22:2, 198-218.

Bakmaz, O., Đuranović-Miličić, J., Dugonjić, D., Dragosavac, M., Dejanović, A., Grublješić, Ž. & Popović, S. (2024-b).Management of non-standard agricultural equipment used in the treatment of patients based on the evaluation of the owners of agricultural holdings and the management of health organizations, Agricultural machinery, 33-38.

Bakmaz, O., Đuranović-Miličić, J., Dragosavac, M., Sredojević, D., Gligović, D., Arnautović, I., Krstajić, G., Pajović, I., & Popović, S. (2025). Financial Management of IT Systems in Medium-Sized Agricultural Enterprises and Healthcare Institutions in the Republic of Serbia. J Agron Technol Eng Manag, 8(4); 1651-1658.

Đajić, D., Đukanović, D., Stanić, S. and Kovačević, K. (2001). Bolesti usta, parodontologija, atlas. Beograd: Elit Medica.

Dessaune, N., Porpino, M.T.M., Antunes, H.d.S., Rodrigues, R.C.V., Perez, A.R., Pires, F.R., Siqueira, J.F., Jr., Armada, L. (2018). Pro-Inflammatory

and Anti-Inflammatory Cytokine Expression in Post-Treatment Apical Periodontitis. J. Appl. Oral Sci., 1-8, 26, e20170455

Dimitrijević, B. (2019). Klinička parodontologija, Zavod za udzbenike, Beograd.

Dragosavac, M., Anđelić, S., Bakmaz, O., Radaković, M., Arnautović, I, Davidov, T., Nastić, S. & Popović, S. (2023). Introducing Software Solutions into the Management of Public Enterprises Whose Founders are Local Self-Government Units in the Example of the Republic of Serbia, Lex Localis – Journal of Local Self-Government, 21(4); 1021–1042.

Jovanović, S., Đokić, S. & Popvić, S. (2025). Procurement of agricultural mechanism through IPARD measure-1, and application of marketing in the meaning of explaining the profitability of investments by the advisory service, Agricultural machinery, 50(2): pp. 1-6.

Lutovac, J., Tomaš-Miskin, S. & Popović, S. (2025). Obtaining loans for energy efficiency depending on the influence of analyzed factors on the decision-making of citizens and small businesses in the Republic of Serbia, International Review (2025No.1-2); 173-181.

Newman, M. and Carranza, F. (2006). Carranza's Clinical Periodontology. 10th Edition, Saunders Co., Lindo.

Newman, M., Takei, H., Klokkevold, P. and Caranzza, F. (2012). Clinical periodontology. Elsevier. St Louis.

Popović, S., Novaković, S., Đuranović, D., Mijić, R., Grublješić, Ž., Aničić, J. & Majstorović, A. (2017). Application of international accounting standard-16 in a public company with predominantly agricultural activities, Economic Research-Ekonomska Istraživanja, 30(1): 1850–1864.

Popović, S., Vitomir, J., Tomaš-Miskin, S., Davidov, T., Nastić., S., Popović, V., Popović, D., Vitomir, G. (2021). The importance of a realistically determined amount of tax on property rights relating to the ownership of agricultural land in the Republic of Serbia adopted by tax authorities of local self-government units, Economics of agriculture, 4: 1029-1042.

Popović, S., Bakmaz, O., Popović, D., Dragosavac, M., Nastić, S., Pajović, I., Majstorović, A., Sredojević, D., Radaković, M. Petković, Z. (2025).

Valuation of agricultural land in relation to location of city centers as a factor in the management of mentioned resource, Agriculture & Forestry, 71(1): 99-113.

Popović, M. (2025). The role of probiotics in therapy periodontopathies-thesis, University of Novi Sad, Faculty of Dentistry.

Radović, M., Vitomir, J. & Popović, S. (2019). The Importance of Implementation of Internal Audit in Enterprises Founded by the Republic of Serbia, Lex Localis - Journal Of Local Self-Government, 17(4): 1001–1011.

Radović, M., Vitomir, J. & Popović, S. (2021-a). Impact of internal control in enterprises founded by local self-government units: the case of Republic of Serbia, Inzinerine Ekonomika-Engineering Economics, 2021, 32(1); 82–90.

Radović, M., Vitomir, J., Popović, S. & Stojanović, A. (2023). The Importance of Establishing Financial Valuation of Fixed Assets in Public Companies who's Founders Are Local Self-Government Units in the Republic of Serbia, Enginering Economics, 34(3); 246-255.

Tomas-Miskin S., Vitomir, J., Popović, S. & Vitomir, G. (2022). Decision-making of Top Management and Internal Audit on the Issue of Archiving Documentation in Companies Founded by Local Government Units in the Republic of Serbia, Lex Localis – Journal of Local Self-Government, 20(4); 889 – 995.

Tošković, J. Popović, S. & Mijić, R. (2016). Was Implementation of transitional reforms in Western Balkans Successful? Actual problems of economics, 11(185); 68-78.

Vihnanek, N. (2015). Application of oral probiotics in periodontology, graduate thesis, Zagreb.

Vitomir, J., Tomaš-Miskin, S., Ivić, M., Popović, S. (2020). Implementation of the Tender by the Municipal Service Administration from the Aspect of Management in the Municipalities of the Republic of Serbia, Lex Localis – Journal of Local Self-Government, 8(3); 469 – 486.

Vitomir, J., Radović, M. & Popović, S. (2021). The Effect of Public Finance Control on the Improvement of Work of Internal Auditors in Enterprises Founded by the Local Self-government Units on the Example of the

Republic of Serbia, Lex localis - Journal of Local Self-Government, 19(2).

Vitomir, J., Tomas-Miskin, S. & Popović, S. (2023). The Importance of the Application of Controlling in the Management of Public Enterprises who's Founders is Local Self-government Units: The Example of Republic of Serbia, Lex Localis – Journal of Local Self-Government, 21(3); 545–565.

CHAPTER 2
FUNDAMENTALS OF ORAL CAVITY BIOLOGY

[1]Prof. Dr. Mehwash KASHIF
[2]Assoc. Prof. Dr. Muhammad Abdullah KAMRAN
[3]Dr. Kainat ARSHAD
[4]Dr. Erum FATIMA

[1]Department of Oral Pathology, Karachi Medical and Dental College, /Karachi Metropolitan University, Pakistan, mehwashkashif@gmail.com, ORCID ID: 0000-0002-3125-8616
[2]Department of Pediatrics and Orthodontic Sciences, College of Dentistry, King Khalid University, Abha, Saudi Arabia mmuhammad@kku.edu.sa, ORCID ID: 0000-0003-2222-0832
[3]Karachi Medical and Dental College,Karachi Metropolitan University, Pakistan, kainatarshad401@gmail.com, ORCID ID: 0009-0001-4915-9077
[4]Karachi Medical and Dental College, /Karachi Metropolitan University, Pakistan, erumfatima192@gmail.com

INTRODUCTION

The oral cavity represents one of the most important anatomical regions of the human body, serving as the initial entry point to both the digestive and respiratory systems. It functions as a critical interface between the external environment and internal physiological systems. From an anatomical standpoint, it is made up of the lips, tongue, teeth, palate, gums, cheeks, and floor of the mouth, each of which has a distinct purpose. The mouth cavity is necessary for speech, chewing, salivation (which activates enzymatic digestion), taste perception, and the initial stages of swallowing. (Akram, 2024). A dynamic biological environment that supports various physiological processes is created by the interaction of these structures. These coordinated functions highlight the complexity and multifunctional nature of the oral cavity. The mouth cavity is a perfect place for a population of bacteria, fungi, protozoa, archaea, and viruses to colonize because it is humid, warm, and nutrient-rich. This community frequently takes the shape of a complex structure known as biofilm or plaque (Abranches et al., 2018). The oral cavity has occasionally been compared to a mirror portraying a person's health (Squier & Kremer, 2001). This analogy emphasizes its diagnostic significance in identifying both local and systemic conditions. The host oral tissues and these bacteria are normally in harmony, but when this equilibrium is disrupted, it can lead to a range of illnesses, including periodontal disease and caries. The connection between dental health and general systemic health is likewise becoming more and more clear. Emerging evidence further suggests associations with cardiovascular diseases, diabetes, and other systemic disorders. Therefore, maintaining oral health and preventing disease require an awareness of the basic principles underlying the mouth cavity. An outline of the basic biological features of the mouth cavity is given in this chapter. This chapter aims to provide a comprehensive understanding that integrates structure, function, and clinical relevance. Its anatomical features, physiological processes, microbial ecology, and roles in preserving oral health are all given special attention.

1. DEVELOPMENT OF THE ORAL CAVITY

The embryologic development of the head and neck is a complex process and is marked by the establishment of the branchial (or pharyngeal) arches (Magreni & May, 2015; Sadler, 2000). Mesoderm, neural crest cells, and ectoderm make up these arches, each of which contributes to particular craniofacial characteristics. The development of the face, palate, and upper oral cavity components is considerably affected by the first pharyngeal arch. The first arch splits into the maxillary and mandibular prominences very early in development. By the end of the fourth week of advancement, these prominences form four separate swellings—two maxillary and two mandibular—and are connected to a superior, midline swelling known as the frontonasal process. The mouth or stomodeum is formed by the depression of the surface ectoderm that is encircled by these five structures (Hiwatashi & Ohno, 2015). Subsequent fusion and growth of these prominences lead to the formation of the upper lip, primary palate, and jaw structures.

2. DEVELOPMENT OF TOOTH

When the oral epithelium thickens and proliferates locally over the developing maxillary and mandibular arches at around 4 weeks, the human embryo shows the earliest indication of tooth development (Sasaki, 1990). This thickening forms the dental lamina, which serves as the origin of all primary teeth. The ectoderm and ectomesenchyme combine to produce the tooth. The ectomesenchyme is primarily derived from neural crest cells, which play a crucial role in craniofacial development. The enamel organ, which is distinct from the primitive oral epithelium lining the stomodeum (primitive oral cavity), is the source of enamel. The structure of the tooth, the development of the tooth's formative cells, and the timing of their secretion are all determined by epithelial-mesenchymal interaction. Tooth development progresses through well-defined stages: bud, cap, and bell, each characterized by specific morphological and cellular changes. The dental papilla develops from mesenchymal cells toward the inner edge of the enamel organ. Additionally, ectomesenchymal cells that are nearer the enamel organ's outer edge develop into dental follicles. Dental papilla is the source of dentine and pulp.

On the other hand, dental follicles are the source of the cementum, periodontal ligament, and alveolar bone.(Kumar, 2023). This coordinated development ensures proper tooth morphogenesis and functional integration within the oral cavity.

3. DEVELOPMENT OF SALIVARY GLANDS

In order to generate and secrete saliva, the mammalian salivary gland grows into a highly branching structure (Patel & Hoffman, 2014). This process, known as branching morphogenesis, is regulated by complex epithelial–mesenchymal signaling pathways. Saliva's primary role is to lubricate the mouth cavity, which is the first stage of food digestion. It also has antibacterial qualities and an ionic composition that encourages tooth mineralization (Chibly et al., 2022). Similar to other ectodermal derivatives like sebaceous and mammary glands, all of the salivary glands in humans originate from the oral cavity's ectoderm. The nasopharyngeal and oral ectoderms give birth to the small salivary glands. They emerge subsequent to the main salivary glands.Human parotid and submandibular gland primordiums emerge in the sixth week of pregnancy, while sublingual gland primordiums emerge after the seventh or eighth week. In the third month, the development of the small salivary glands starts (Kumar, 2023). The maturation of these glands involves lumen formation, acinar differentiation, and establishment of ductal systems essential for saliva secretion.

4. ANATOMICAL ORGANIZATION OF THE ORAL CAVITY

The first part of the digestive system is the oral cavity, also referred to as the mouth or buccal cavity. It is comprised up of various anatomically distinct components that collaborate well to carry out a variety of tasks. Further, it plays a crucial role in both protective and sensory processes by acting as a transitional area between the gastrointestinal system and the outside world. The vestibule, which is the space between the cheeks, teeth, and lips, and the oral cavity proper are the two distinct areas that make up the oral cavity, which is encircled by the lips (Kamrani & Sadiq, Updated 2023 Aug 14).

These areas have specific organs and tissues that work together to support biological and mechanical functions. The oral cavity proper lies internal to the dental arches and communicates posteriorly with the oropharynx through the oropharyngeal isthmus. Studying both normal function and abnormal circumstances requires an understanding of its anatomical arrangement.

5. BOUNDARIES OF THE ORAL CAVITY

Clearly defined anatomical limits enclose the interior structures of the mouth cavity. The lips form the anterior boundary of the oral cavity, the cheeks provide the lateral boundary, the hard palate forms the superior boundary, and the mucosa lining the tongue's upper surface and the sheet of muscles that attach to the inner side of the mandible—the geniohyoid, mylohyoid, and digastric—form the inferior boundary. The posterior boundary is marked by the oropharyngeal isthmus, which connects the oral cavity to the pharynx and plays a role in swallowing. The upper and lower dentition, the tongue, salivary glands, mucosal glands, and the mucosal tissue covering the hard palate—which contains the rugae are among the structures found in this area, despite the fact that it is essentially a passageway for food and air. Together, the connective tissue lamina propria and the specific oral epithelium that lines these structures offer protection and sensory abilities. The pharyngeal cavity, a more intricate and rather uneven area, is continuous with the oral cavity (German & Palmer, 2006). These boundaries help maintain the structural integrity and functional efficiency of the oral cavity.

6. MAJOR COMPONENT AND BIOLOGICAL IMPORTANCE

The mouth cavity has a number of important parts that support its biological and functional activities. The primary purpose of teeth, which are essential organs for vertebrates, is to bite and chew food into little bits. Human teeth are always a major factor in look and beauty, and they are crucial for everything from protecting support organs to pronouncing words correctly (Chen & Liu, 2014). Additionally, teeth play a part in occlusal stability and help preserve vertical face dimensions. Another essential part is the tongue.

Normal breathing, swallowing, oral transport, emesis, coughing, and, in humans, speech production all depend on the tongue (Sokoloff & Burkholder, 2013). Also it has specific sensory receptors for taste perception and uses lingual tonsils to support immunological defense. The oral mucosa is classified as either keratinized or nonkeratinized, and its histology differs from place to site. With the exception of the tongue dorsum, hard palatal mucosa, and connected gingiva, the majority of oral mucosa is nonkeratinized (Woo, 2024). This variation reflects adaptation to functional demands such as mastication, protection, and flexibility. Human salivary glands (SG) are essential for preserving the homeostasis of the mouth cavity. They produce and release saliva, a multipurpose fluid that protects the oral mucosa and tooth surface by providing mucosal lubrication, salivary electrolytes, antibacterial chemicals, and several enzymes (Carpenter, 2013; de Paula et al., 2017; Feller et al., 2013). Saliva also plays a crucial role in buffering oral pH and initiating digestion through enzymes such as amylase. Gingiva is the collective word for another significant tissue found inside the oral cavity. They fulfill two purposes. They are a part of both the periodontium and the oral mucous membrane. Clinically speaking, the gingiva, also known as the marginal periodontium, is a collar of masticating mucosa that surrounds the teeth of the entire deciduous or permanent dentition and is connected to both the teeth and the alveolar process. It is composed of both connective and epithelial tissues (Schroeder & Listgarten, 1997). It also serves as a protective barrier against mechanical stress and microbial invasion.

All things considered, the mouth cavity's anatomical arrangement shows a highly coordinated system in which structural elements are intimately linked to functional tasks. This organization supports defense mechanisms and the preservation of oral and systemic health in addition to facilitating vital functions like mastication, speaking, and swallowing. Clinical diagnosis and therapeutic approaches in oral healthcare both depend on an understanding of its arrangement.

7. ORAL MUCOSA STRUCTURE AND FUNCTION

The oral mucosa is the moist, soft tissue membrane that lines the structures of the oral cavity. It extends anteriorly from the junction of the vermilion border of the lips and the labial mucosa to posteriorly reach the palatopharyngeal folds, serving as a protective and functional interface within the mouth.

Classification of Oral Mucosa:

The oral mucosa can be classified into three types as follows.

- Lining (Movable) Mucosa: The lining mucosa covers the mobile regions of the oral cavity, including the soft palate, cheeks, lips, alveolar mucosa, floor of the mouth, and vestibular fornix. It is characterized by a non-keratinized stratified squamous epithelium (Groeger & Meyle, 2019), which provides flexibility and resilience to accommodate movements during speech, mastication, and facial expressions.
- Masticatory mucosa: Masticatory mucosa refers to the firm, immobile mucosa that is tightly bound to the underlying bone, found in the attached gingiva and hard palate. It is covered by keratinized or parakeratinized stratified squamous epithelium, which enables it to withstand the mechanical stresses of mastication and maintain a stable attachment to the bone.
- Specialized mucosa: Specialized mucosa is located on the dorsum of the tongue and is composed of stratified squamous epithelium, which may be keratinized or non-keratinized. It is termed “specialized” due to the presence of various types of lingual papillae and taste buds, which are essential for the perception of taste (Garg et al., 2025), Because the dorsum of the tongue actively participates in mastication, this mucosa is sometimes also classified as masticatory mucosa.

Histological Structure of Oral Mucosa:

Histologically, the oral mucosa is formed by three layers

- Oral Epithelium: The oral epithelium is a stratified squamous surface layer, with its thickness and degree of keratinization varying according to its location and the functional demands placed upon it (Groeger & Meyle, 2019).

- Lamina Propria: Beneath the oral epithelium lies the lamina propria, a connective tissue layer that provides structural support and nutrition to the overlying epithelium.

Composition: The lamina propria is primarily composed of fibrous connective tissue, consisting of a network of type I and III collagen fibers along with elastin in certain regions. Fibroblasts are the predominant cells, responsible for producing and maintaining the extracellular matrix and fibers.

Layers of Lamina Propria:

The lamina propria is organized into two distinct layers:

- Papillary Layer: The papillary layer is the superficial portion of the lamina propria. It is composed of loose connective tissue within the dermal papillae and contains blood vessels and nerve fibers.
- Dense Layer: The dense layer forms the deeper portion of the lamina propria and consists of dense connective tissue with a high concentration of fibers. Between the papillary and dense layers lies a capillary plexus, which supplies nutrients to all layers of the mucosa and sends capillaries into the papillary layer (Williams et al., 2021).
- Submucosa: The submucosa is a layer of dense, irregular connective tissue located at the deepest level of the oral mucosa. In certain regions, it is absent, and the lamina propria attaches directly to underlying bone or muscle. The submucosa is typically present in movable areas such as the cheeks and lips, providing flexibility and cushioning. In contrast, in regions like the attached gingiva and hard palate, the oral mucosa is firmly bound to the periosteum of the underlying bone, with no intervening submucosa (Iglesias-Sancho et al., 2020). Submucosa contains fat, minor salivary glands, main blood vessels, and nerves supplying the mucosa. It contains adipose tissue, minor salivary glands, major blood vessels, and nerves supplying the mucosa.

Functions of Oral Mucosa:

- Protective Function: The oral epithelium serves as a critical barrier against both physiological and pathogenic stresses. It provides physical and immune protection, preventing the invasion of harmful microorganisms.

- This defense is achieved through multiple layers of epithelial cells connected by tight cell–cell junctions, along with immune surveillance mediated by dendritic cells (DCs) and T helper 17 (Th17) cells, that regulate local immune responses.(Wang et al., 2025).
- Secretion: The oral mucosa primarily contributes to secretion through saliva, produced by the major and minor salivary glands, whose ducts open into the oral cavity maintaining moisture and lubrication. Sebaceous glands may also be present in labial and buccal mucosa, producing lipid-rich secretions with a possible protective role.(Wertz, 2018).
- Sensory Function: The oral cavity receives sensory innervation primarily from the three branches of the trigeminal nerve. The oral mucosa contains three main types of sensory receptors—Merkel's discs, Meissner's corpuscles, and free nerve endings—which enable the perception of temperature, touch, and pain. It also contributes to taste perception, including the five basic tastes: salty, sweet, sour, bitter, and umami, with emerging evidence suggesting sensitivity to fat. (Garg et al., 2025). The taste receptor cells are located primarily on the dorsum of the tongue and soft palate.
- Healing Capacity: The oral mucosa tends to heal faster and with less scar formation compared to the skin due to rich vascular supply, rapid epithelial turnover.

Biology of teeth:

Teeth are extremely specialised mineralised structures that are essential to the oral cavity. Mastication, which is the mechanical disintegration of food into smaller pieces, is their main job. They also contribute to speech, aesthetics, and maintenance of occlusal relationships. In terms of biology, teeth are made up of both organic and inorganic materials. Hydroxyapatite crystals, which offer strength and hardness, make up the majority of the inorganic component. Collagen and other proteins that support structure and durability are part of the organic component. Three distinct hard tissues enamel, dentin, and cementum as well as the soft pulp tissue make up a tooth (Kamakura, 2015). Together, these tissues guarantee the tooth's longevity, sensitivity, and nutrition.

Studying the function and pathology of teeth requires an understanding of their biological structure and cellular makeup.

Tooth Structure: A tooth's structure is made up of several specialized tissues that are placed in a very orderly fashion. A tooth is divided into two main parts one is above gum line known as crown and other is bellow gum line known as root as shown in Fig 1. The crown is clinically visible, while the root is embedded within the alveolar bone and provides anchorage. White enamel covers the crown and shields the teeth. Despite being the strongest material in the body, enamel has very little capacity to heal itself when it is harmed. Dentin, which resembles bone but is tougher, lies beneath the enamel. It forms the bulk of the tooth and provides structural support to enamel. The core (pulp) chamber, which houses connective tissue, blood vessels, and nerves, is surrounded by dentin. Dentin is sensitive to temperature fluctuations and touch (Sulyanto, Modified Jul 2025). The firm, avascular connective tissue that surrounds tooth roots is called cementum ("Chapter 9 - Periodontium," 2013). It facilitates attachment of periodontal ligament fibers to the root surface. It serves as a barrier to prevent bacterial evasion and limit epithelial development (Arzate et al., 2015; Wang et al., 2025). These structures work together to give the tooth resilience, nutrition, and sensory function. Damage to any of these components can compromise tooth integrity and lead to pathological conditions.



Figure 1. This illustration demonstrates the anatomical structure of a tooth, highlighting the crown and root regions along with enamel, dentin, and pulp cavity.

8. CELLULAR COMPONENTS OF TEETH

Specialised cells are essential for the development and upkeep of dental structures. The characteristics of tooth enamel are ultimately determined by ameloblasts, the cells responsible for its formation (Hubbard, 1998). Enamel is produced during tooth development through a highly regulated process of biomineralization. However, its capacity for regeneration is extremely limited due to the loss of ameloblasts following tooth eruption. Particular kinds of cells called odontoblasts create dentin and have distinctive morphological traits, such as extending cytoplasmic extensions into dentinal tubules. While mesenchymal stem cells produce osteoblasts, which are typical hard-tissue-forming cells, during normal and pathological bone metabolism, odontoblasts primarily form dentin, but there is growing evidence that they also function as nociceptors and defensive cells in the dental pulp (Kawashima & Okiji, 2016). The organic matrix of cementum is secreted by specialized cells called cementoblasts, which create cementum. Cellular cementum is produced in some areas when these cells become stuck in the matrix and develop into cementocytes, while acellular cementum is produced in other areas when embedded cells are absent. Cementum formation also involves two types of fibers. Extrinsic (Sharpey's) fibers, which represent the embedded ends of principal periodontal ligament fibers, are primarily produced by fibroblasts, with partial contribution from cementoblasts. In contrast, intrinsic fibers, which constitute the fibers of cementum proper, are secreted exclusively by cementoblasts (Yamamoto et al., 2016). Additionally, immunological cells found in the pulp aid in the body's defense against infection. These include macrophages, dendritic cells, and lymphocytes that contribute to pulpal immunity. These cellular elements are essential for the growth, maintenance, and defensive processes of teeth.

9. CLINICAL RELEVANCE OF TOOTH BIOLOGY

Effective diagnosis and treatment of dental problems depend on an understanding of tooth biology. The significance of early prevention and restorative care in caries management is explained by knowledge of enamel's limited potential for regeneration. Pain perception in diseases like pulpitis and dentin hypersensitivity is closely correlated with the sensitivity of dentin and the existence of odontoblastic processes.

The importance of cementum in periodontal health and illness is highlighted by its function in periodontal ligament attachment. Furthermore, immune cells in the pulp both contribute to inflammatory reactions and aid in defense against infection, thus prompt management is essential to maintaining tooth life and function.

9.1 Salivary Gland and Salivary Biology

Reduced salivary flow, or hyposalivation, can have serious effects on both oral and overall health. It may lead to a diminished sense of taste and cause difficulties in eating, swallowing, and speaking. In the oral cavity, it increases the risk of tooth decay, disrupts the natural balance of oral microorganisms, and contributes to the development of oral lesions, infections, gum disease, and periodontitis. Beyond oral health, hyposalivation can negatively impact general well-being and may even result in nutritional deficiencies due to challenges in food intake (Azuma et al., 2018; Bergdahl & Bergdahl, 2000; Lan et al., 2020). These effects collectively reduce quality of life.

9.2 Salivary Gland and Secretions

The three major pairs of salivary glands in humans—the parotid glands (PGs), submandibular glands (SMGs), and sublingual glands (SLGs)—are responsible for producing over 90% of the approximately 1 to 1.5 liters of saliva secreted daily. The remainder is produced by minor salivary glands distributed throughout the oral mucosa (Patel & Hoffman, 2014).

9.3 Composition of Saliva

Human saliva consists of roughly 99% water and 1% a mixture of organic and inorganic substances. This 1% includes:

Proteins and enzymes that aid digestion and protect oral tissues, Immunological components that help defend against pathogens, and electrolytes such as calcium, phosphate, and sodium (Carpenter, 2013). These components collectively contribute to its multifunctional role in oral health.

9.4 Functions of Saliva

Saliva plays a major role in maintaining the oral environment and keeping the oral cavity healthy. Some major functions of saliva are well described in Table 1.

Table 1. Major Physiological Functions of Saliva and Their Key Bioactive Components

Function	Description	Key Components	References
Lubrication & Digestion	Facilitates chewing, swallowing, and speech; begins carbohydrate digestion	Water, mucins, amylase	(Carpenter, 2013)
Antimicrobial Protection	Inhibits microbial growth and maintains oral microbiome balance	Immunoglobulins, lysozyme, lactoferrin	(Carpenter, 2013)
Remineralization	Repairs early enamel demineralization	Calcium, phosphate ions	
Wound Healing	Promotes rapid healing of oral tissues	EGF, FGF, VEGF, histatins,	(Bodner & Dayan, 1995; Brand & Veerman, 2013; T et al., 2019; Xu et al., 2019)

Diagnostic Approach: Saliva serves as a valuable diagnostic fluid because its molecular composition mirrors the body's physiological and pathological conditions. It contains biomarkers such as proteins, nucleic acids, and hormones derived from local and systemic sources (Gardner et al., 2020; Ishikawa et al., 2016; Yoshizawa et al., 2013; Zhang et al., 2012). By integrating salivary transcriptomics with proteomics, researchers can determine the specific contributions of the major salivary glands as well as serum components to the overall salivary composition (Saitou et al., 2020). Saliva is increasingly used as a non-invasive alternative to blood tests.

Saliva has shown promise as a source of biomarkers for a wide range of conditions, including dental caries, periodontal disease, cancer, diabetes, viral infections, Sjögren's syndrome, cardiovascular disease, and numerous other systemic disorders (Zhang et al., 2016). This potential for saliva to be used as a tool in the diagnosis of disease has been widely recognized and reviewed previously (Kaufman & Lamster, 2002; Pereira et al., 2020).

Role in Transmission: Saliva also plays a role in viral transmission and diagnosis. Certain viruses, including SARS-CoV-2, can replicate in salivary gland epithelium and be detected in saliva, supporting its role in disease transmission and monitoring (Baghizadeh Fini, 2020; Chen et al., 2020; Han & Ivanovski, 2020).

Physiology of Salivary Secretion: Salivary glands are exocrine organs responsible for saliva production and secretion. The submandibular glands (SMGs) primarily contribute to basal saliva flow during rest, whereas the parotid glands (PGs), the largest salivary glands, produce most of the stimulated saliva. This secretion is typically serous and amylase-rich. Salivary secretion is tightly regulated by the autonomic nervous system, through coordinated sympathetic and parasympathetic activity.

Anatomy of the Salivary Gland: The exocrine organs known as salivary glands are in charge of producing and secreting saliva, which is essential for preserving oral homeostasis. Based on their size and contribution to the overall amount of saliva produced, they are generally divided into major and small salivary glands. The parotid, submandibular, and sublingual glands are the three main salivary glands; each has a different location, ductal system, and secretion type. Table 2 summarizes these glands' primary anatomical characteristics.

Table 2. This Table Summarizes Primary Anatomical Characteristics of Salivary Glands.

Gland	Location	Duct	Secretion Type	References
Parotid	Near ear, posterior to mandible	Stensen's duct (opens near upper 2nd molar)	Mainly serous	(Frommer, 1977; Kessler & Bhatt, 2018; Yakirevich et al., 2010)
Submandibular	Below mandible, digastric region	Wharton's duct (opens at sublingual caruncle)	Mixed (serous dominant)	(Ellis, 2012; Kessler & Bhatt, 2018)
Sublingual	Floor of mouth, submucosal	Multiple small ducts	Mainly mucous	(de Paula et al., 2017)

Histology of Salivary Glands: Salivary glands (SGs) are highly branched exocrine organs organized into a complex ductal network that terminates in secretory units called acini. Acini consist of serous and mucous cells producing watery and viscous secretions, respectively. Saliva is modified through intercalated, striated, and excretory ducts before secretion. Myoepithelial cells (MECs) surround the acini and ducts, contracting in response to neuronal signals to aid in saliva expulsion. The glandular microenvironment includes ECM, autonomic innervation, immune cells, and vascular supply supporting function (de Paula et al., 2017).

Saliva, produced by major and minor salivary glands, is essential for oral and systemic health. It supports digestion, lubrication, antimicrobial defense, wound healing, and remineralization, while also serving as a non-invasive diagnostic tool.

10. ORAL MICROBIOME

The human oral cavity harbors a diverse and complex microbial community, collectively known as the oral microbiome (Deo & Deshmukh, 2019)

It has diverse microorganisms, including bacteria,archaea, fungi, and viruses (Wade, 2013) existing predominantly as biofilms that maintain oral homeostasis by fostering a balanced microbial environment (Deo & Deshmukh, 2019). A balanced oral microbiome serves as a natural defense barrier, inhibiting colonization by pathogenic species and preventing oral infections (Kilian et al., 2016). It also helps prevent oral diseases such as dental caries and periodontal diseases. (Deo & Deshmukh, 2019; Kilian et al., 2016).

Under healthy conditions, the oral microbiome exists in a balanced, commensal relationship with its environment, similar to microbial communities in the skin, gut, or vagina. However, certain opportunistic microorganisms can shift to a pathogenic state under specific conditions, contributing to the onset of both oral and systemic diseases (Zarco et al., 2012).

Major oral diseases, including dental caries and periodontal disease, both result from and contribute to imbalances in the oral microbiota, a condition known as dysbiosis. This disruption may also contribute to systemic conditions such as cardiovascular disease and inflammatory bowel disease (Thomas et al., 2021).

10.1 Composition and Functional Role of the Oral Microbiome

With over 700 known bacterial species, the oral microbiome is the second most diverse microbiota within the human body (Deo & Deshmukh, 2019). It includes bacterial phyla, including Actinobacteria, Bacteroidetes, Chlamydia, Euryarchaeota, Fusobacteria, Firmicutes, Proteobacteria, Spirochaetes, and Tenericutes(Dewhirst et al., 2010).

In a healthy oral cavity, the bacterial community includes both Gram-positive and Gram-negative species. Among the Gram-positive cocci, genera such as Abiotrophia, Peptostreptococcus, Streptococcus, and Stomatococcus play key roles. Some species, including Streptococcus sanguinis and Streptococcus salivarius, produce antimicrobial compounds that help maintain oral health (Huang et al., 2011).

10.2 Oral Microbiome Dysbiosis

Oral microbiome dysbiosis refers to an imbalance or disruption in the composition and functioning of the oral microbial community (Deo & Deshmukh, 2019). Factors such as poor oral hygiene, diet, medical conditions, medications, and lifestyle can disrupt the microbiome. This imbalance alters the harmonious interactions among microbial species, affecting overall microbial ecology. Oral dysbiosis is associated with common conditions such as dental caries, periodontal disease, halitosis, and oral candidiasis, and has also been linked to systemic disorders, including cardiovascular disease, diabetes (Thomas et al., 2021). A key characteristic of oral microbiome dysbiosis is the overgrowth of harmful bacteria and a decline in beneficial microorganisms (DeGruttola et al., 2016). This promotes inflammation and tissue damage.

10.3 Oral Disease Associates with Microbiome Dysbiosis

Dental Caries

Dental caries is a common infectious disease driven by bacterial pathogens and characterized by progressive dental tissue destruction (Bowen et al., 2018). Key cariogenic bacteria include Prevotella spp., Lactobacillus spp., Dialister microorganisms and Filifactor spp. have surfaced as potential contributors to caries pathogenesis and advancement (Tanner et al., 2018). Caries develop due to acidogenic bacteria that metabolize sugars, producing acids that cause enamel demineralization (Selwitz et al., 2007).

Gingivitis

Gingivitis is inflammation of the gums caused by plaque accumulation due to poor oral hygiene. Plaque accumulates in the gingival sulcus, increasing infection risk. Key species include Porphyromonas gingivalis, Streptococcus spp, Aggregatibacter actinomycetemcomitans, and Fusobacterium spp (JO & P, 2023). If left untreated, gingivitis can progress to periodontitis, a more severe condition. Thus, good oral hygiene is essential (Intan Suhana et al., 2020).

Periodontitis

The oral microbiome is central to the development and progression of periodontal disease, which can advance from gingivitis to deeper destruction of the supporting periodontal tissues (Gao et al., 2018; Lu et al., 2019). Dental plaque forms on both supragingival and subgingival surfaces, promoting the growth of microbial biofilms on teeth. The gingival crevice and crevicular epithelium serve as key sites for microbial colonization, aided by gingival crevicular fluid and the anaerobic conditions of the sulcus (Malla et al., 2019)..Subgingival plaque, which contains over 500 microbial species, undergoes significant compositional changes as periodontal disease progresses (Moore & Moore, 1994). Under healthy conditions, Gram-positive cocci and rods dominate as early colonizers, while Fusobacterium nucleatum acts as a bridging species facilitating bacterial coaggregation (Moore & Moore, 1994). As gingivitis progresses, Gram-negative bacteria, including Prevotella spp. and Fusobacterium nucleatum ssp., become more prevalent, correlating with increased levels of pro-inflammatory cytokines such as IL-1, IL-6, IL-17, TNF-α, and IL-23 (Curtis et al., 2020; Gao et al., 2018).

Oral Thrush

Oral thrush, or oral candidiasis, is a fungal infection of the oral mucosa caused by the overgrowth of Candida albicans (Singh et al., 2014). Other species include Candida glabrata and Candida tropicalis (Rajendra Santosh et al., 2021). Symptoms include white patches and mucosal inflammation. In severe cases, it may lead to pain and a burning sensation in the mouth (J & EH, 2026). Risk factors for oral thrush include poor oral hygiene, a weakened immune system, and medical conditions such as diabetes or HIV/AIDS (Rajendra Santosh et al., 2021).

11. ORAL HEALTH AND DENTAL CARE PRACTICES

Probiotics

Dental probiotics help maintain oral microbiota balance by colonizing the oral mucosa, producing bioactive metabolites that support ecological equilibrium, and enhancing the host's immune defenses (Kamel et al., 2009).

Common probiotic strains include Streptococcus, Lactobacillus, and Bifidobacterium species. These inhibit pathogenic bacteria and reduce inflammation.

A balanced oral microbiome is essential for oral health. Dysbiosis leads to diseases such as caries, gingivitis, periodontitis, and candidiasis, with possible systemic effects. Preventive care and microbial balance are key to maintaining oral and systemic health.

12. IMMUNE DEFENSE MECHANISM IN THE ORAL CAVITY

12.1 Saliva and Oral Epithelial Barrier

Saliva plays a fundamental role in maintaining oral homeostasis by lubricating oral surfaces and facilitating mechanical clearance of microorganisms. The oral mucosa, composed primarily of stratified squamous epithelium, serves as both a physical and immunological barrier against microbial invasion. Continuous epithelial cell desquamation contributes significantly to host defense by removing adherent microorganisms from oral surfaces, with exfoliated epithelial cells constituting approximately 55–60% of total salivary cellular content (Choudhury et al., 2020; Theda et al., 2018).

Epithelial cell shedding is a key defense mechanism, preventing microbial colonization and removing adherent bacteria—each cell may clear up to 100 organisms from oral surfaces (Dawes, 2003). Detached, pathogen-bound cells are swallowed and neutralized by gastric acid and salivary antimicrobial components, (Rao et al., 2006) highlighting saliva's role in mechanical pathogen clearance and oral-systemic protection

12.2 Cellular Components of Salivary Immunity

Epithelial Cells

Epithelial cells are the most abundant cellular components in saliva and act as the first line of defense by preventing microbial adhesion and facilitating pathogen removal through desquamation.

Neutrophils (Polymorphonuclear Leukocytes)

Polymorphonuclear neutrophils (PMNs) are the second most prevalent cellular component in saliva and constitute the primary leukocyte population within the oral environment. They comprise approximately 30–40% of the total salivary cell count and account for nearly 95–98% of all leukocytes present in saliva (Choudhury et al., 2020; Vidović et al., 2012).

Neutrophils play a pivotal role in the innate immune system, through phagocytosis, degranulation, reactive oxygen species production, and neutrophil extracellular trap formation, enabling rapid control of microbial invasion.

Oral Dendritic Cells (Dcs)

Dendritic cells act as antigen-presenting cells linking innate and adaptive immunity by capturing microbial antigens and activating T lymphocytes in regional lymph nodes.

12.3 Adaptive Immune Response in Saliva

Lymphocytes (T and B cells)

Lymphocytes are responsible for specific and long-lasting immune responses. (Alberts et al., 2007). Upon infection, antigen-presenting cells (APCs), activated by the innate immune system, migrate to lymph nodes to initiate adaptive responses (Angel et al., 2009). In lymph nodes, naïve T cells recognize presented antigens, differentiate into $CD4^+$ or $CD8^+$ T cells, and return to infection sites to mediate immunity. In the oral cavity, $CD8^+$ T cells contribute to inflammation regulation and may reduce bone loss by inhibiting osteoclastogenesis (Cardoso & Arosa, 2017; Choi et al., 2001).

12.4 Salivary Immune Molecules

Secretory Immunoglobulin A (SIgA)

Secretory IgA is the predominant salivary immunoglobulin and plays a major role in mucosal defense by inhibiting microbial adhesion and neutralizing pathogens.

Salivary Cytokines

Salivary cytokines regulate inflammatory responses and contribute to host defense. Dysregulation of cytokine production is associated with oral inflammatory diseases such as gingivitis and periodontitis and may reflect systemic disease activity such as HIV, tuberculosis, and oral head and neck cancer (Diesch et al., 2021)

Lactoferrin

Lactoferrin is an antimicrobial glycoprotein with antibacterial and antiviral activity secreted by exocrine glands and neutrophils that contributes to innate immune protection within saliva.

Mucin

Mucins are salivary glycoproteins that exhibit a gel-like consistency, form a protective barrier over oral surfaces, inhibit microbial colonization, exhibit direct antimicrobial activity, and participate in cell signaling processes (Frenkel & Ribbeck, 2015).

Salivary mucins, including MUC5B, MUC7, MUC19, MUC1, and MUC4. MUC7 specifically contains binding sites for bacteria such as Streptococcus sanguinis, S. sobrinus, and S. oralis (Frenkel & Ribbeck, 2015).

13. GINGIVAL BARRIER AND GINGIVAL CREVICULAR FLUID

Gingival crevicular fluid is an inflammatory exudate presented within the gingival sulcus that transports immune cells, cytokines, enzymes, and microbial products. It plays an important role in maintaining periodontal homeostasis and reflects the inflammatory status of periodontal tissues.

Oral Cavity and Systemic Health

According to the WHO (World Health Organization), approximately 3.5 billion individuals are affected by oral diseases, making them one of the most prevalent non-communicable disease globally. The oral cavity serves as the primary entry point to the human body. Under normal conditions, the oral microbiota exists in a balanced state and contributes to maintaining oral health.

However, when systemic health is compromised or oral hygiene is inadequate, these normally harmless microorganisms can become pathogenic, leading to both local and systemic infections (Lalla & Papapanou, 2011).

Oral Systemic Connection

Historically, oral bacteria were considered to have effects limited to the oral cavity. However, growing evidence has established a strong link between oral health and systemic diseases, with 200 possible connections between systemic diseases and oral health reported by The American Dental Association (Kapila, 2021; Loos, 2006; Nazir et al., 2019; Peres et al., 2019; Watt et al., 2019).

The detrimental effects of oral infections on overall health are largely attributed to the entry of oral microorganisms or their byproducts into the bloodstream, a process known as bacteremia. Dental caries, primarily caused by Streptococcus Mutans and Lactobacillus species, exemplify such oral disorders. Oral pathogens can spread to distant organs through local oral circulation or via the gastrointestinal tract into systemic circulation. Upon reaching target organs, these microbes can alter immune responses and trigger the release of inflammatory mediators, contributing to the development of systemic diseases (Arweiler & Netuschil, 2016; Kitamoto et al., 2020; Peng et al., 2022; Varoni & Rimondini, 2022) such as Atrial Fibrillation (AF) (Park et al., 2023), CVS (Loos, 2006), Liver Cirrhosis (Pitones-Rubio et al., 2020) (Chan, 2024), Inflammatory bowel disease (IBD)(Park et al., 2021).

Oral Health and Major Systemic Disease

Recent studies indicate that oral health can influence multiple organ systems. Effects range from metabolic disturbances, such as insulin dysregulation associated with periodontal disease, to more complex systemic complications contributing to neurodegenerative disorders (King et al., 2022) even an adverse pregnancy outcome due to bacteremia by releasing prostaglandins which may stimulate oxytocin, promoting preterm labor and low birth weight.

Cardiovascular Risk

A 14-year study found that periodontal disease patients had a

25% higher risk of developing CVD than those who are orally healthy counterparts. (Dhadse et al., 2010). Men younger than 50 years with periodontal disease demonstrate 72% more risk of developing CVD. Mechanistically, Oral pathogens, specifically Streptococcus mutans (cariogenic) and Porphyromonas gingivalis (perioodontitis), induce platelet aggregation, leading to thrombus formation (Nomura et al., 2014). Furthermore, 42% of atheromas are found in periodontal disease patients. (Reddy et al., 2000). Patients who are susceptible to endocarditis may need prophylactic antibiotics prior to invasive oral surgical procedures.

Hypertension

Chronic oral infections, causing bacteremia trigger the release of pro-inflammatory cytokines such as tumor necrosis factor-alpha (TNF-α) and interleukin-6 (IL-6). These cytokines sustain local inflammation and, when circulated systemically, can adversely affect vascular tissues (Yildirim et al., 2022). Elevated levels of pro-inflammatory cytokines impair endothelial function, reducing the production of nitric oxide (NO), a key mediator of vasodilation. This results in increased arterial stiffness and diminished vascular flexibility, which raises vascular resistance and contributes to the onset of hypertension (Ja'far, 2017; Nomura et al., 2020).

Diabetes

The diabetes and oral health connection is a multifaceted and bidirectional relationship (Borgnakke, 2019). Evidence indicates that regular tooth brushing linked to a reduced risk of developing diabetes (Chang et al., 2020). Diabetes can cause increased susceptibility to infection, impaired immune response and dry mouth (xerostomia) (Borgnakke & Poudel, 2021).

Dry mouth can increase the risk of dental caries and periodontal issues. On the other hands Diabetes can weaken the immune system, reducing the body’s ability to combat oral infections. Conversely, oral infections can allow pathogenic bacteria to enter the bloodstream, heightening the risk of complications in individuals with diabetes (Nasseh et al., 2014).

Neurodegenerative Disorder

Alzheimer's Disease

Oral health is linked to neurodegenerative disorders such as Alzheimer's disease, as chronic periodontal inflammation may promote neurodegeneration through systemic inflammatory mediators, increased blood–brain barrier permeability, and invasion of pathogens or their toxins from inflamed periodontal tissues. Gums contribute to Alzheimer's disease (Sedghi et al., 2021).

Oral health is strongly linked to systemic health, as poor oral hygiene and chronic oral infections are associated with conditions such as cardiovascular disease, diabetes, adverse pregnancy outcomes, neurodegenerative disorders, hypertension, and certain cancers. These effects occur through systemic inflammation, bacteremia, immune dysregulation, and the spread of oral pathogens, highlighting the importance of maintaining good oral hygiene and regular dental care.

Key Takeaways

The oral cavity is a highly specialized and dynamic biological system that plays a central role in both local and systemic health. As the initial point of contact between the external environment and the human body, it performs essential functions including mastication, speech, taste perception, and the initiation of digestion. These functions are supported by a complex integration of anatomical structures such as the teeth, tongue, oral mucosa, and salivary glands, each contributing uniquely to the overall physiology of the oral environment.

From a developmental perspective, the formation of oral and craniofacial structures is a precisely regulated process involving intricate epithelial–mesenchymal interactions. This coordinated development ensures the proper formation of teeth, salivary glands, and supporting tissues, all of which are essential for functional efficiency and long-term oral health.

At the microscopic level, the oral cavity is characterized by specialized tissues and cellular components that provide protection, sensation, and repair. The oral mucosa acts as a critical barrier against mechanical and microbial challenges, while also facilitating rapid healing.

Similarly, the structural and cellular complexity of teeth enables them to perform mechanical functions while maintaining sensitivity and defense against external stimuli.

Saliva plays a fundamental role in maintaining oral homeostasis through its multifunctional properties, including lubrication, digestion, antimicrobial defense, remineralization, and wound healing. In addition, its emerging role as a non-invasive diagnostic fluid highlights its importance beyond the oral cavity, offering valuable insights into systemic health.

A key feature of the oral environment is the presence of a diverse and complex microbiome. Under normal conditions, this microbial community exists in a balanced state, contributing to host defense and maintaining ecological stability. However, disruption of this balance, known as dysbiosis, can lead to the development of common oral diseases such as dental caries, gingivitis, periodontitis, and oral candidiasis. These conditions not only affect oral tissues but may also have broader implications for systemic health.

The oral cavity is also equipped with a sophisticated immune defense system involving both innate and adaptive mechanisms. Saliva, epithelial barriers, immune cells, and antimicrobial molecules work in coordination to prevent microbial invasion and maintain tissue integrity. This defense system is essential in preserving the delicate balance between host and microorganisms.

Importantly, increasing evidence highlights a strong relationship between oral health and systemic diseases. Oral infections and inflammation can contribute to conditions such as cardiovascular disease, diabetes, hypertension, and neurodegenerative disorders through mechanisms including bacteremia, chronic inflammation, and immune dysregulation. This oral–systemic connection underscores the significance of oral health as an integral component of overall well-being.

Conclusion:

In conclusion, the oral cavity should not be viewed in isolation but as an integral part of the human body, where structure, function, microbiology, and immunity are closely interconnected. A thorough understanding of these fundamental aspects is essential for the prevention, diagnosis, and management of oral diseases.

For undergraduate students, this foundation provides a critical framework for advancing clinical knowledge and emphasizes the importance of maintaining oral health as a key determinant of general health.

REFERENCES

Abranches, J., Zeng, L., Kajfasz, J. K., Palmer, S. R., Chakraborty, B., Wen, Z. T., Richards, V. P., Brady, L. J., & Lemos, J. A. (2018). Biology of Oral Streptococci. Microbiology Spectrum, 6(5), 10.1128/microbiolspec.gpp1123-0042-2018.

Akram, M. (2024). Oral Cavity. Journal of Dental and Oral Care, 3(1).

Alberts, B., Johnson, A., Lewis, J., Raff, M., Roberts, K., & Walter, P. (2007). Molecular Biology of the Cell (5th, Ed.). W. W. Norton & Company. https://doi.org/https://doi.org/10.1201/9780203833445

Angel, C. E., Chen, C.-J. J., Horlacher, O. C., Winkler, S., John, T., Browning, J., MacGregor, D., Cebon, J., & Dunbar, P. R. (2009). Distinctive localization of antigen-presenting cells in human lymph nodes. Blood, The Journal of the American Society of Hematology, 113(6), 1257-1267.

Arweiler, N. B., & Netuschil, L. (2016). The Oral Microbiota. Adv Exp Med Biol, 902, 45-60. https://doi.org/10.1007/978-3-319-31248-4_4

Arzate, H., Zeichner-David, M., & Mercado-Celis, G. (2015). Cementum proteins: role in cementogenesis, biomineralization, periodontium formation and regeneration. Periodontology 2000, 67(1), 211-233.

Azuma, N., Katada, Y., & Sano, H. (2018). Deterioration in saliva quality in patients with Sjögren's syndrome: impact of decrease in salivary epidermal growth factor on the severity of intraoral manifestations. Inflammation and Regeneration, 38(1), 6. https://doi.org/10.1186/s41232-018-0062-0

Baghizadeh Fini, M. (2020). Oral saliva and COVID-19. Oral Oncology, 108, 104821. https://doi.org/https://doi.org/10.1016/j.oraloncology.2020.104821

Bergdahl, M., & Bergdahl, J. (2000). Low unstimulated salivary flow and subjective oral dryness: association with medication, anxiety, depression, and stress. J Dent Res, 79(9), 1652-1658. https://doi.org/10.1177/00220345000790090301

Bodner, L., & Dayan, D. (1995). Epithelium and connective tissue regeneration during palatal wound healing in desalivated rats--a comparative study. Comp Biochem Physiol A Physiol, 111(3), 415-419. https://doi.org/10.1016/0300-9629(95)00036-7

Borgnakke, W. S. (2019). IDF Diabetes Atlas: Diabetes and oral health – A two-way relationship of clinical importance. Diabetes Research and Clinical Practice, 157. https://doi.org/10.1016/j.diabres.2019.107839

Borgnakke, W. S., & Poudel, P. (2021). Diabetes and Oral Health: Summary of Current Scientific Evidence for Why Transdisciplinary Collaboration Is Needed [Perspective]. Frontiers in Dental Medicine, Volume 2 - 2021. https://doi.org/10.3389/fdmed.2021.709831

Bowen, W. H., Burne, R. A., Wu, H., & Koo, H. (2018). Oral Biofilms: Pathogens, Matrix, and Polymicrobial Interactions in Microenvironments. Trends Microbiol, 26(3), 229-242. https://doi.org/10.1016/j.tim.2017.09.008

Brand, H. S., & Veerman, E. C. (2013). Saliva and wound healing. Chin J Dent Res, 16(1), 7-12.

Cardoso, E. M., & Arosa, F. A. (2017). CD8+ T Cells in Chronic Periodontitis: Roles and Rules [Opinion]. Frontiers in Immunology, Volume 8 - 2017. https://doi.org/10.3389/fimmu.2017.00145

Carpenter, G. H. (2013). The secretion, components, and properties of saliva. Annual review of food science and technology, 4(1), 267-276.

Chan, W. S. H. (2024). The role of oral health in the prevention of systemic diseases. Universal Library of Medical and Health Sciences, 1(1).

Chang, Y., Lee, J. S., Lee, K. J., Woo, H. G., & Song, T. J. (2020). Improved oral hygiene is associated with decreased risk of new-onset diabetes: a nationwide population-based cohort study. Diabetologia, 63(5), 924-933. https://doi.org/10.1007/s00125-020-05112-9

Chapter 9 - Periodontium. (2013). In A. Nanci (Ed.), Ten Cate's Oral Histology (Eighth Edition) (pp. 205-232). Mosby. https://doi.org/https://doi.org/10.1016/B978-0-323-07846-7.00009-4

Chen, H., & Liu, Y. (2014). Chapter 2 - Teeth. In J. Z. Shen & T. Kosmač (Eds.), Advanced Ceramics for Dentistry (pp. 5-21). Butterworth-Heinemann. https://doi.org/https://doi.org/10.1016/B978-0-12-394619-5.00002-X

Chen, L., Zhao, J., Peng, J., Li, X., Deng, X., Geng, Z., Shen, Z., Guo, F., Zhang, Q., Jin, Y., Wang, L., & Wang, S. (2020). Detection of SARS-

CoV-2 in saliva and characterization of oral symptoms in COVID-19 patients. Cell Prolif, 53(12), e12923. https://doi.org/10.1111/cpr.12923

Chibly, A. M., Aure, M. H., Patel, V. N., & Hoffman, M. P. (2022). Salivary gland function, development, and regeneration. Physiological Reviews, 102(3), 1495-1552. https://doi.org/10.1152/physrev.00015.2021

Choi, Y., Woo, K. M., Ko, S. H., Lee, Y. J., Park, S. J., Kim, H. M., & Kwon, B. S. (2001). Osteoclastogenesis is enhanced by activated B cells but suppressed by activated CD8(+) T cells. Eur J Immunol, 31(7), 2179-2188. https://doi.org/10.1002/1521-4141(200107)31:7<2179::aid-immu2179>3.0.co;2-x

Choudhury, S. N., Novotny, M., Aevermann, B. D., Lee, S., Mandava, A., Qian, Y., & Freire, M. (2020). A protocol for revealing oral neutrophil heterogeneity by single-cell immune profiling in human saliva.

Curtis, M. A., Diaz, P. I., & Van Dyke, T. E. (2020). The role of the microbiota in periodontal disease. Periodontology 2000, 83(1), 14-25. https://doi.org/https://doi.org/10.1111/prd.12296

Dawes, C. (2003). Estimates, from salivary analyses, of the turnover time of the oral mucosal epithelium in humans and the number of bacteria in an edentulous mouth. Arch Oral Biol, 48(5), 329-336. https://doi.org/10.1016/s0003-9969(03)00014-1

de Paula, F., Teshima, T. H. N., Hsieh, R., Souza, M. M., Nico, M. M. S., & Lourenco, S. V. (2017). Overview of Human Salivary Glands: Highlights of Morphology and Developing Processes. The Anatomical Record, 300(7), 1180-1188. https://doi.org/https://doi.org/10.1002/ar.23569

DeGruttola, A. K., Low, D., Mizoguchi, A., & Mizoguchi, E. (2016). Current Understanding of Dysbiosis in Disease in Human and Animal Models. Inflamm Bowel Dis, 22(5), 1137-1150. https://doi.org/10.1097/mib.0000000000000750

Deo, P. N., & Deshmukh, R. (2019). Oral microbiome: Unveiling the fundamentals. J Oral Maxillofac Pathol, 23(1), 122-128. https://doi.org/10.4103/jomfp.JOMFP_304_18

Dewhirst, F. E., Chen, T., Izard, J., Paster, B. J., Tanner, A. C. R., Yu, W.-H., Lakshmanan, A., & Wade, W. G. (2010). The Human Oral Microbiome.

Journal of Bacteriology, 192(19), 5002-5017. https://doi.org/doi:10.1128/jb.00542-10

Dhadse, P., Gattani, D., & Mishra, R. (2010). The link between periodontal disease and cardiovascular disease: How far we have come in last two decades ? J Indian Soc Periodontol, 14(3), 148-154. https://doi.org/10.4103/0972-124x.75908

Diesch, T., Filippi, C., Fritschi, N., Filippi, A., & Ritz, N. (2021). Cytokines in saliva as biomarkers of oral and systemic oncological or infectious diseases: A systematic review. Cytokine, 143, 155506. https://doi.org/https://doi.org/10.1016/j.cyto.2021.155506

Ellis, H. (2012). Anatomy of the salivary glands. Surgery (Oxford), 30(11), 569-572. https://doi.org/https://doi.org/10.1016/j.mpsur.2012.09.008

Feller, L., Altini, M., Khammissa, R., Chandran, R., Bouckaert, M., & Lemmer, J. (2013). Oral mucosal immunity. Oral surgery, oral medicine, oral pathology and oral radiology, 116(5), 576-583.

Frenkel, E. S., & Ribbeck, K. (2015). Salivary mucins in host defense and disease prevention. J Oral Microbiol, 7, 29759. https://doi.org/10.3402/jom.v7.29759

Frommer, J. (1977). The human accessory parotid gland: its incidence, nature, and significance. Oral Surg Oral Med Oral Pathol, 43(5), 671-676. https://doi.org/10.1016/0030-4220(77)90049-4

Gao, L., Xu, T., Huang, G., Jiang, S., Gu, Y., & Chen, F. (2018). Oral microbiomes: more and more importance in oral cavity and whole body. Protein & Cell, 9(5), 488-500. https://doi.org/10.1007/s13238-018-0548-1

Gardner, A., Carpenter, G., & So, P. W. (2020). Salivary Metabolomics: From Diagnostic Biomarker Discovery to Investigating Biological Function. Metabolites, 10(2). https://doi.org/10.3390/metabo10020047

Garg, R., Trillo, G., & Brizuela, M. (2025). Anatomy, Head and Neck, Tongue Taste Buds. In StatPearls [Internet]. StatPearls Publishing.

German, R. Z., & Palmer, J. B. (2006). Anatomy and development of oral cavity and pharynx. GI Motility online.

Groeger, S., & Meyle, J. (2019). Oral Mucosal Epithelial Cells [Review]. Frontiers in Immunology, Volume 10 - 2019. https://doi.org/10.3389/fimmu.2019.00208

Han, P., & Ivanovski, S. (2020). Saliva—Friend and Foe in the COVID-19 Outbreak. Diagnostics, 10(5), 290. https://www.mdpi.com/2075-4418/10/5/290

Hiwatashi, N., & Ohno, S. (2015). Development and Regeneration of the Oral Cavity and the Pharynx. In J. Ito (Ed.), Regenerative Medicine in Otolaryngology (pp. 109-127). Springer Japan. https://doi.org/10.1007/978-4-431-54856-0_8

Huang, R., Li, M., & Gregory, R. L. (2011). Bacterial interactions in dental biofilm. Virulence, 2(5), 435-444. https://doi.org/10.4161/viru.2.5.16140

Hubbard, M. J. (1998). Enamel Cell Biology Towards a Comprehensive Biochemical Understanding. Connective Tissue Research, 38(1-4), 17-32. https://doi.org/10.3109/03008209809017013

Iglesias-Sancho, M., Llambí Mateos, F., & Salleras-Redonnet, M. (2020). Drug Compounding for Diseases of the Oral Mucosa. Actas Dermo-Sifiliográficas (English Edition), 111(10), 822-828. https://doi.org/https://doi.org/10.1016/j.adengl.2020.08.013

Intan Suhana, M. A., Farha, A., & Hassan, B. M. (2020). Inflammation of the Gums. Malays Fam Physician, 15(1), 71-73.

Ishikawa, S., Sugimoto, M., Kitabatake, K., Sugano, A., Nakamura, M., Kaneko, M., Ota, S., Hiwatari, K., Enomoto, A., Soga, T., Tomita, M., & Iino, M. (2016). Identification of salivary metabolomic biomarkers for oral cancer screening. Scientific Reports, 6(1), 31520. https://doi.org/10.1038/srep31520

J, J., & EH, T. (2026). Candidiasis. In S. Publishing (Ed.), StatPearls. StatPearls Publishing. https://www.ncbi.nlm.nih.gov/books/NBK560624/

Ja'far, Z. J. a. (2017). Dental Caries Experience and Severity in Relation to Salivary TNF- ? and AlP in a Group of Overweight Children Compared with Normal Weight Children. International Journal of Science and Research (IJSR), 6(8), 2061-2065. https://doi.org/https://dx.doi.org/10.21275/ART20173902

JO, D., & P, J. (2023). Gingivitis. In S. Publishing (Ed.), StatPearls. StatPearls Publishing. https://www.ncbi.nlm.nih.gov/books/NBK557422/

Kamakura, S. (2015). Tooth and Tooth-Supporting Structures. In M. Niinomi, T. Narushima, & M. Nakai (Eds.), Advances in Metallic Biomaterials: Tissues, Materials and Biological Reactions (pp. 99-122). Springer Berlin Heidelberg. https://doi.org/10.1007/978-3-662-46836-4_5

Kamel, U. F., Maddison, P., & Whitaker, R. (2009). Impact of primary Sjögren's syndrome on smell and taste: effect on quality of life. Rheumatology, 48(12), 1512-1514. https://doi.org/10.1093/rheumatology/kep249

Kamrani, P., & Sadiq, N. M. (Updated 2023 Aug 14). Anatomy, Head and Neck, Oral Cavity (Mouth). In StatPearls. StatPearls Publishing 2026 Jan. https://www.ncbi.nlm.nih.gov/books/NBK545271/

Kapila, Y. L. (2021). Oral health's inextricable connection to systemic health: Special populations bring to bear multimodal relationships and factors connecting periodontal disease to systemic diseases and conditions. Periodontol 2000, 87(1), 11-16. https://doi.org/10.1111/prd.12398

Kaufman, E., & Lamster, I. B. (2002). The diagnostic applications of saliva--a review. Crit Rev Oral Biol Med, 13(2), 197-212. https://doi.org/10.1177/154411130201300209

Kawashima, N., & Okiji, T. (2016). Odontoblasts: Specialized hard-tissue-forming cells in the dentin-pulp complex. Congenital Anomalies, 56(4), 144-153. https://doi.org/https://doi.org/10.1111/cga.12169

Kessler, A. T., & Bhatt, A. A. (2018). Review of the Major and Minor Salivary Glands, Part 1: Anatomy, Infectious, and Inflammatory Processes. J Clin Imaging Sci, 8, 47. https://doi.org/10.4103/jcis.JCIS_45_18

Kilian, M., Chapple, I. L. C., Hannig, M., Marsh, P. D., Meuric, V., Pedersen, A. M. L., Tonetti, M. S., Wade, W. G., & Zaura, E. (2016). The oral microbiome – an update for oral healthcare professionals. British Dental Journal, 221(10), 657-666. https://doi.org/10.1038/sj.bdj.2016.865

King, S., Chow, C. K., & Eberhard, J. (2022). Oral health and cardiometabolic disease: understanding the relationship. Intern Med J, 52(2), 198-205. https://doi.org/10.1111/imj.15685

Kitamoto, S., Nagao-Kitamoto, H., Hein, R., Schmidt, T. M., & Kamada, N. (2020). The Bacterial Connection between the Oral Cavity and the Gut

Diseases. Journal of Dental Research, 99(9), 1021-1029. https://doi.org/10.1177/0022034520924633

Kumar, G. (2023). Orban's Oral Histology & Embryology-E-BOOK. Elsevier Health Sciences.

Lalla, E., & Papapanou, P. N. (2011). Diabetes mellitus and periodontitis: a tale of two common interrelated diseases. Nat Rev Endocrinol, 7(12), 738-748. https://doi.org/10.1038/nrendo.2011.106

Lan, X., Chan, J. Y. K., Pu, J. J., Qiao, W., Pang, S., Yang, W. F., Wong, K. C. W., Kwong, D. L. W., & Su, Y. X. (2020). Saliva electrolyte analysis and xerostomia-related quality of life in nasopharyngeal carcinoma patients following intensity-modulated radiation therapy. Radiother Oncol, 150, 97-103. https://doi.org/10.1016/j.radonc.2020.06.016

Loos, B. (2006). Systemic effects of periodontitis. Annals of the Royal Australasian College of Dental Surgeons, 18, 27-29.

Lu, M., Xuan, S., & Wang, Z. (2019). Oral microbiota: A new view of body health. Food Science and Human Wellness, 8(1), 8-15. https://doi.org/https://doi.org/10.1016/j.fshw.2018.12.001

Magreni, A., & May, J. G. (2015). Embryology of the oral structures. Operative Techniques in Otolaryngology-Head and Neck Surgery, 26(3), 110-114. https://doi.org/https://doi.org/10.1016/j.otot.2015.06.002

Malla, M. A., Dubey, A., Kumar, A., Yadav, S., Hashem, A., & Abd_Allah, E. F. (2019). Exploring the Human Microbiome: The Potential Future Role of Next-Generation Sequencing in Disease Diagnosis and Treatment [Review]. Frontiers in Immunology, Volume 9 - 2018. https://doi.org/10.3389/fimmu.2018.02868

Moore, W. E. C., & Moore, L. V. H. (1994). The bacteria of periodontal diseases. Periodontology 2000, 5(1), 66-77. https://doi.org/https://doi.org/10.1111/j.1600-0757.1994.tb00019.x

Nasseh, K., Greenberg, B., Vujicic, M., & Glick, M. (2014). The Effect of Chairside Chronic Disease Screenings by Oral Health Professionals on Health Care Costs. American Journal of Public Health, 104(4), 744-750. https://doi.org/10.2105/ajph.2013.301644

Nazir, M. A., Izhar, F., Akhtar, K., & Almas, K. (2019). Dentists' awareness about the link between oral and systemic health. J Family Community Med, 26(3), 206-212. https://doi.org/10.4103/jfcm.JFCM_55_19

Nomura, R., Matayoshi, S., Otsugu, M., Kitamura, T., Teramoto, N., & Nakano, K. (2020). Contribution of Severe Dental Caries Induced by Streptococcus mutans to the Pathogenicity of Infective Endocarditis. Infection and Immunity, 88(7), 10.1128/iai.00897-00819. https://doi.org/doi:10.1128/iai.00897-19

Nomura, R., Otsugu, M., Naka, S., Teramoto, N., Kojima, A., Muranaka, Y., Matsumoto-Nakano, M., Ooshima, T., & Nakano, K. (2014). Contribution of the Interaction of Streptococcus mutans Serotype k Strains with Fibrinogen to the Pathogenicity of Infective Endocarditis. Infection and Immunity, 82(12), 5223-5234. https://doi.org/doi:10.1128/iai.02164-14

Park, J. H., Lee, H., Kim, J. W., & Song, T. J. (2023). Association between periodontal disease status and risk of atrial fibrillation: a nationwide population-based cohort study. BMC Oral Health, 23(1), 461. https://doi.org/10.1186/s12903-023-03165-x

Park, S.-Y., Hwang, B.-O., Lim, M., Ok, S.-H., Lee, S.-K., Chun, K.-S., Park, K.-K., Hu, Y., Chung, W.-Y., & Song, N.-Y. (2021). Oral–Gut Microbiome Axis in Gastrointestinal Disease and Cancer. Cancers, 13(9), 2124. https://www.mdpi.com/2072-6694/13/9/2124

Patel, V. N., & Hoffman, M. P. (2014). Salivary gland development: a template for regeneration. Semin Cell Dev Biol, 25-26, 52-60. https://doi.org/10.1016/j.semcdb.2013.12.001

Peng, X., Cheng, L., You, Y., Tang, C., Ren, B., Li, Y., Xu, X., & Zhou, X. (2022). Oral microbiota in human systematic diseases. International Journal of Oral Science, 14(1), 14. https://doi.org/10.1038/s41368-022-00163-7

Pereira, J. A. M., Porto-Figueira, P., Taware, R., Sukul, P., Rapole, S., & Câmara, J. S. (2020). Unravelling the Potential of Salivary Volatile Metabolites in Oral Diseases. A Review. Molecules, 25(13), 3098. https://www.mdpi.com/1420-3049/25/13/3098

Peres, M. A., Macpherson, L. M. D., Weyant, R. J., Daly, B., Venturelli, R., Mathur, M. R., Listl, S., Celeste, R. K., Guarnizo-Herreño, C. C., Kearns, C., Benzian, H., Allison, P., & Watt, R. G. (2019). Oral diseases: a global public health challenge. Lancet, 394(10194), 249-260. https://doi.org/10.1016/s0140-6736(19)31146-8

Pitones-Rubio, V., Chávez-Cortez, E. G., Hurtado-Camarena, A., González-Rascón, A., & Serafín-Higuera, N. (2020). Is periodontal disease a risk factor for severe COVID-19 illness? Med Hypotheses, 144, 109969. https://doi.org/10.1016/j.mehy.2020.109969

Rajendra Santosh, A. B., Muddana, K., & Bakki, S. R. (2021). Fungal Infections of Oral Cavity: Diagnosis, Management, and Association with COVID-19. SN Comprehensive Clinical Medicine, 3(6), 1373-1384. https://doi.org/10.1007/s42399-021-00873-9

Rao, A., Jump, R. L., Pultz, N. J., Pultz, M. J., & Donskey, C. J. (2006). In vitro killing of nosocomial pathogens by acid and acidified nitrite. Antimicrob Agents Chemother, 50(11), 3901-3904. https://doi.org/10.1128/aac.01506-05

Reddy, M. S., Geurs, N. C., Jeffcoat, R. L., Proskin, H., & Jeffcoat, M. K. (2000). Periodontal disease progression. Journal of periodontology, 71(10), 1583-1590.

Sadler, T. (2000). Head and neck. Langman's medical embryology, 345-381.

Saitou, M., Gaylord, E. A., Xu, E., May, A. J., Neznanova, L., Nathan, S., Grawe, A., Chang, J., Ryan, W., Ruhl, S., Knox, S. M., & Gokcumen, O. (2020). Functional Specialization of Human Salivary Glands and Origins of Proteins Intrinsic to Human Saliva. Cell Rep, 33(7), 108402. https://doi.org/10.1016/j.celrep.2020.108402

Sasaki, T. (1990). Cell biology of tooth enamel formation: functional electron microscopic monographs.

Schroeder, H. E., & Listgarten, M. A. (1997). The gingival tissues: the architecture of periodontal protection. Periodontology 2000, 13(1), 91-120. https://doi.org/https://doi.org/10.1111/j.1600-0757.1997.tb00097.x

Sedghi, L. M., Bacino, M., & Kapila, Y. L. (2021). Periodontal Disease: The Good, The Bad, and The Unknown [Review]. Frontiers in Cellular and Infection Microbiology, Volume 11 - 2021.

Selwitz, R. H., Ismail, A. I., & Pitts, N. B. (2007). Dental caries. The Lancet, 369(9555), 51-59. https://doi.org/10.1016/S0140-6736(07)60031-2

Singh, A., Verma, R., Murari, A., & Agrawal, A. (2014). Oral candidiasis: An overview. J Oral Maxillofac Pathol, 18(Suppl 1), S81-85. https://doi.org/10.4103/0973-029x.141325

Sokoloff, A., & Burkholder, T. (2013). Tongue Structure and Function. In L. K. McLoon & F. Andrade (Eds.), Craniofacial Muscles: A New Framework for Understanding the Effector Side of Craniofacial Muscle Control (pp. 207-227). Springer New York. https://doi.org/10.1007/978-1-4614-4466-4_12

Squier, C. A., & Kremer, M. J. (2001). Biology of Oral Mucosa and Esophagus. JNCI Monographs, 2001(29), 7-15.

Sulyanto, R. (Modified Jul 2025). Biology of the Teeth. Merck & Co., Inc. Retrieved March 18 from https://www.msdmanuals.com/home/mouth-and-dental-disorders/biology-of-the-mouth-and-teeth/biology-of-the-teeth

T, V., AM, R., AS, S., & MA, J.-R. (2019). The power of saliva: Antimicrobial and beyond. PLoS Pathogens, 15(11), e1008058.

Tanner, A. C. R., Kressirer, C. A., Rothmiller, S., Johansson, I., & Chalmers, N. I. (2018). The Caries Microbiome: Implications for Reversing Dysbiosis. Advances in Dental Research, 29(1), 78-85.

Theda, C., Hwang, S. H., Czajko, A., Loke, Y. J., Leong, P., & Craig, J. M. (2018). Quantitation of the cellular content of saliva and buccal swab samples. Scientific Reports, 8(1), 6944. https://doi.org/10.1038/s41598-018-25311-0

Thomas, C., Minty, M., Vinel, A., Canceill, T., Loubières, P., Burcelin, R., Kaddech, M., Blasco-Baque, V., & Laurencin-Dalicieux, S. (2021). Oral Microbiota: A Major Player in the Diagnosis of Systemic Diseases. Diagnostics, 11(8), 1376. https://www.mdpi.com/2075-4418/11/8/1376

Varoni, E. M., & Rimondini, L. (2022). Oral Microbiome, Oral Health and Systemic Health: A Multidirectional Link. Biomedicines, 10(1), 186. https://www.mdpi.com/2227-9059/10/1/186

Vidović, A., Vidović Juras, D., Vučićević Boras, V., Lukač, J., Grubišić-Ilić, M., Rak, D., & Sabioncello, A. (2012). Determination of leucocyte

subsets in human saliva by flow cytometry. Arch Oral Biol, 57(5), 577-583. https://doi.org/10.1016/j.archoralbio.2011.10.015

Wade, W. G. (2013). The oral microbiome in health and disease. Pharmacological Research, 69(1), 137-143.

Wang, H., Yang, L., Huang, X., Ma, L., Wang, C., Wang, X., & Cao, Z. (2025). Cementum regeneration strategies: Insights from development and periodontal microenvironment [Review]. Frontiers in Cell and Developmental Biology, Volume 13 - 2025.

Watt, R. G., Daly, B., Allison, P., Macpherson, L. M. D., Venturelli, R., Listl, S., Weyant, R. J., Mathur, M. R., Guarnizo-Herreño, C. C., Celeste, R. K., Peres, M. A., Kearns, C., & Benzian, H. (2019). Ending the neglect of global oral health: time for radical action. Lancet, 394(10194), 261-272. https://doi.org/10.1016/s0140-6736(19)31133-x

Wertz, P. W. (2018). Lipids and the Permeability and Antimicrobial Barriers of the Skin. Journal of Lipids, 2018(1), 5954034. https://doi.org/https://doi.org/10.1155/2018/5954034

Williams, D. W., Greenwell-Wild, T., Brenchley, L., Dutzan, N., Overmiller, A., Sawaya, A. P., Webb, S., Martin, D., Hajishengallis, G., Divaris, K., Morasso, M., Haniffa, M., & Moutsopoulos, N. M. (2021). Human oral mucosa cell atlas reveals a stromal-neutrophil axis regulating tissue immunity. Cell, 184(15), 4090-4104.e4015. https://doi.org/10.1016/j.cell.2021.05.013

Woo, S.-B. (2024). 1 - Introduction. In S.-B. Woo (Ed.), Oral Pathology (Third Edition) (pp. 1-9). Elsevier. https://doi.org/https://doi.org/10.1016/B978-0-323-82918-2.00010-X

Xu, F., Laguna, L., & Sarkar, A. (2019). Aging-related changes in quantity and quality of saliva: Where do we stand in our understanding? Journal of Texture Studies, 50(1), 27-35.

Yakirevich, E., Sabo, E., Klorin, G., Alos, L., Cardesa, A., Ellis, G. L., Shumway, B. S., & Gnepp, D. R. (2010). Primary mucin-producing tumours of the salivary glands: a clinicopathological and morphometric study. Histopathology, 57(3), 395-409. https://doi.org/10.1111/j.1365-2559.2010.03639.x

Yamamoto, T., Hasegawa, T., Yamamoto, T., Hongo, H., & Amizuka, N. (2016). Histology of human cementum: Its structure, function, and development. Jpn Dent Sci Rev, 52(3), 63-74.

Yildirim, B. G., Aksit, C., Mutlu, M., Ainola, M., Eklund, K. K., Leskelä, J., Pussinen, P., & Beklen, A. (2022). Severity and progression rate of periodontitis are associated with an increased risk of hypertension of patients attending a university clinic. BMC Oral Health, 22(1), 627. https://doi.org/10.1186/s12903-022-02637-w

Yoshizawa, J. M., Schafer, C. A., Schafer, J. J., Farrell, J. J., Paster, B. J., & Wong, D. T. (2013). Salivary biomarkers: toward future clinical and diagnostic utilities. Clin Microbiol Rev, 26(4), 781-791. https://doi.org/10.1128/cmr.00021-13

Zarco, M., Vess, T., & Ginsburg, G. (2012). The oral microbiome in health and disease and the potential impact on personalized dental medicine. Oral Diseases, 18(2), 109-120. https://doi.org/https://doi.org/10.1111/j.1601-0825.2011.01851.x

Zhang, A., Sun, H., & Wang, X. (2012). Saliva metabolomics opens door to biomarker discovery, disease diagnosis, and treatment. Appl Biochem Biotechnol, 168(6), 1718-1727. https://doi.org/10.1007/s12010-012-9891-5

Zhang, C. Z., Cheng, X. Q., Li, J. Y., Zhang, P., Yi, P., Xu, X., & Zhou, X. D. (2016). Saliva in the diagnosis of diseases. Int J Oral Sci, 8(3), 133-137. https://doi.org/10.1038/ijos.2016.38

CHAPTER 3
INTERNET OF THINGS (IOT) IN MEDICAL SCIENCES: APPLICATIONS, CHALLENGES, AND FUTURE PERSPECTIVES

[1]Dr. Maqubool HOSAIN
[2]Dr. Naveen TRIVEDI
[3]Prof. Dr.Vijaya LAXMI

[1]BIT Mesra, Ranchi, Jharkhand, India, mhosain@bitmesra.ac.in, ORCID ID: 0000-0001-6340-2179
[2]BIT Mesra, Ranchi, Jharkhand, India, ntrivedi@bitmesra.ac.in, ORCID ID: 0000-0003-3843-9409
[3]BIT Mesra, Ranchi, Jharkhand, India, vlaxmi@bitmesra.ac.in, ORCID ID: 0000-0003-0709-5834

INTRODUCTION

The Internet of Things (IoT) is defined as a system consisting of interrelated computing devices, mechanical and digital machines, objects, people, or any other items with embedded electronics, software, sensors, actuators, and connectivity which can connect and exchange data over the internet. These devices not only interact with one another but can also interact with a central health care system to automatically collect and transmit data and make decisions in real time. In the context of the healthcare industry, IoT technology holds immense promise as one of the most disruptive technologies available today.

Healthcare systems across the world are undergoing rapid transformation due to several major challenges, including increasing patient demand, growing aging populations, shortage of healthcare professionals, and continuously rising treatment costs. Chronic diseases such as diabetes, hypertension, cardiovascular disorders, and respiratory illnesses require regular monitoring and long-term management, placing significant pressure on hospitals and clinics. Traditional healthcare models are often reactive in nature, where treatment begins only after symptoms become severe or emergencies occur. This approach leads to higher healthcare expenses, delayed diagnosis, overcrowded hospitals, and reduced quality of care. IoT technology helps overcome these limitations by shifting healthcare from reactive treatment to proactive and preventive care.

Modern-day health care systems are experiencing significant transformation because of numerous factors that have been emerging over time. Some of the most common issues include increasing patients' needs for medical care, aging population, lack of health professionals, and the constant increase in the cost of treatments. Diseases such as diabetes, high blood pressure, heart-related conditions, and lung ailments necessitate routine examination and treatment, which causes a considerable burden on hospital services. Conventional health care systems are mostly reactive, implying that treatments can only begin once the condition reaches its peak or becomes an emergency. As a result, medical facilities experience increased expenses, late diagnosis, overcrowding, and poor quality of service.

The Internet of Things also finds its application in hospital administration and clinical processes. In intelligent hospitals, IoT-based devices facilitate the tracking of assets and inventory, managing patients' flow, regulating the temperature of medicines, and maintaining critical medical equipment. For instance, IoT devices can be used to track the movements of wheelchairs, ventilators, infusion pumps, and other hospital equipment. Smart beds can help monitor the movement of patients, identify any fall incidents, and change bed settings based on their comfort preferences.

Another key strength of the Internet of Things in the field of medical sciences is that it offers a lot of assistance in preventing diseases as well as identifying symptoms at an early stage. With continuous monitoring and collection of health data, IoT solutions can help detect any irregularities and give prior warnings to both doctors and patients about the possibility of developing any serious disease. For managing health crises caused by infectious diseases, IoT temperature scanning systems will play a critical role.

Combining IoT with other emerging technologies like AI, cloud computing, big data, and 5G communication networks makes it more effective than before. The use of AI in analyzing patients' information to predict health issues and suggest solutions will be of immense benefit to people. Cloud technology helps in storing the information in a safer place and easily accessing patients' information through a secure connection. 5G communication network also ensures fast and reliable connections.

Although the technology offers several advantages, it still faces certain difficulties like the risk of data breaches, cyber attacks, compatibility issues, compliance problems, and expensive implementation processes. Nevertheless, through constant advancements and enhanced security protocols, these obstacles can be effectively addressed. To conclude, IoT is revolutionizing the healthcare industry through interconnected and intelligent healthcare systems. The increasing popularity of IoT technology is expected to bring about significant changes to the future of global medicine.

1. ARCHITECTURE OF IOT IN HEALTHCARE

1.1 Sensing Layer

The sensing layer is the basic and initial layer of the architecture of the Internet of Things. The task of this layer is to gather data from the physical environment using different types of sensors, RFIDs, cameras, GPS units, and smart gadgets. They detect and sense real-world parameters, including temperature, humidity, pressure, motion, light level, heart rate, and geolocation. Next, the sensed information is converted to digital form for processing by the IoT's next layers. In the health sector, the layer performs an important function when using wearable gadgets, such as smart thermometers, glucometers, heart-rate detectors, and ECGs. The importance of this layer lies in its responsibility for receiving the highest-quality data. Hence, it is necessary for developing effective IoT solutions.

Includes ECG sensors, BP monitors, SpO_2 sensors, temperature sensors.



Figure 1. Sensing Layer

1.2 Network Layer

The network layer is one of the important parts of IoT architecture, which allows transferring data from sensory devices to other layers in the system hierarchy. In medical sciences and smart healthcare systems, the network layer serves for the transfer of data generated by sensory devices to the processors, cloud, and applications of the health system. Various types of devices including ECG sensors, blood pressure sensors, pulse oximeters, glucose sensors, temperature sensors, and others work constantly, generating patient data that have to be transmitted in an appropriate way to other layers of the architecture. Communication technologies like Bluetooth, ZigBee, Wi-Fi, RFID, NFC, Ethernet, 4G, 5G networks, and others help communicate between devices using various communication channels.

This layer guarantees successful transmission of information to ensure that doctors receive patient data in an accurate manner without any delays. When using remote patient monitoring devices, the network layer helps patients remain in their homes while their health-related information is sent straight to the hospital or healthcare providers. Telemedicine applications benefit from this layer due to easy video conferencing capabilities. Security measures like encryption and authentication become necessary in this layer due to the need for safeguarding patient-related data against any kind of security breach.

Includes Wi-Fi, Bluetooth, Zigbee, and 5G.



Figure 2. Network Layer

1.3 Processing Layer

The processing layer is an essential component of any IoT architecture and plays a significant role in the medical sciences field and other intelligent healthcare systems. The processing layer involves collecting data coming from the sensing layer via the network layer and translating it into useful information. Essentially, the processing layer works as a 'brain' of any IoT system by processing and managing healthcare data generated by numerous sources within that system. In a medical IoT ecosystem, various sensors such as ECG monitor, blood pressure sensor, temperature sensor, and SpO_2 continuously produce patient's data. The processing layer includes servers, cloud computing facilities, databases, and other data management techniques. However, it may also incorporate advanced technologies such as Artificial Intelligence (AI), Machine Learning (ML), and Big Data analytics in order to find abnormalities, predict diseases and assist in making clinical decisions. If the heart rate of a particular patient becomes abnormal, the processing layer processes that information almost instantly and provides necessary alerts to physicians or caretakers. The processing layer stores patients' records securely and provides access to these data whenever it is necessary. Thus, the processing layer significantly improves the accuracy, efficiency, and speed of healthcare services.

Includes cloud computing, edge computing, AI/ML.



Figure 3. Network Layer

1.4 Application Layer

Layer Application is the highest level of the IoT architecture and acts as an interface between the IoT architecture and end-users. In the field of medical science, this layer provides healthcare information in an understandable and usable form for doctors, nurses, patients, and hospital management staff. It takes the output from the processing layer and presents it as valuable services to the end-user in the form of health monitoring dashboard, mobile application, emergency alarm, notification, report, and telemedicine. Using the Application Layer, doctors can monitor a patient's heartbeat, blood pressure, oxygen levels, body temperature, and other parameters from a remote location.

The application layer facilitates better communication between the healthcare provider and the patient by offering video conferencing, appointment scheduling, and prescriptions online. This layer aids in decision-making through the provision of accurate data through reports, graphs, and even historic medical records. The system is able to send out immediate warnings to either the doctor or relative if unusual results arise in an emergency case. There are various security measures offered by this layer as well to ensure that sensitive information is not disclosed.

Includes dashboards, mobile apps, tele-medicine systems.



Figure 4. Layered architecture of IoT-based healthcare system.

2. APPLICATIONS OF IOT IN MEDICAL SCIENCES

2.1 Remote Patient Monitoring

The Remote Patient Monitoring (RPM) is a sophisticated IoT application in the field of healthcare, which enables doctors and other healthcare professionals to keep track of their patients remotely using smart medical equipment. RPM is especially beneficial for elderly patients, people with long-term illnesses, and individuals undergoing post-surgical recovery. In such systems, connected devices like blood pressure monitoring units, glucose monitoring units, electrocardiogram (ECG) sensors, pulse oximeters, temperature sensors, and even activity trackers gather data regarding the well-being of patients constantly. The collected data is then transferred through the internet and/or wireless connections to servers and cloud computing services managed by the healthcare organizations. Physicians can view the gathered data in real-time using dashboard interfaces and mobile apps and act accordingly if any abnormality occurs. For instance, when there is a sudden decrease in oxygen saturation levels or blood pressure levels, immediate alerts can be triggered for the concerned parties.

Remote patient monitoring minimizes unnecessary trips to hospitals, saves money in healthcare expenses, and promotes patient convenience. Additionally, it motivates patients to take an active role in their health management process. In the case of pandemics and emergency situations, RPM was extremely useful for monitoring patients from the comfort of their homes. RPM has greatly improved early detection, proper treatment, and constant care. The integration of IoT technologies, telecommunications, and data analysis in RPM has revolutionized healthcare.

Continuous monitoring reduces hospital visits and improves outcomes.

2.2 Wearable Devices

Wearable technology refers to the use of smart electronic devices which are worn by people for the purpose of monitoring their health, fitness, and activities in real time. They are one of the major applications of the IoT technology in healthcare and medical science. They include items such as smartwatches, fitness bracelets, smart rings, chest bands, and other medical wearables which have sensors.

Such devices can track various parameters including heart rate, blood pressure, body temperature, oxygen saturation level, quality and quantity of sleep, steps taken, calories burnt, and physical activities performed by individuals.

Doctors and other health care professionals can leverage this data to keep track of their patients without having to physically see them, which helps them identify health issues in their patients in their early stages. The benefits of wearable technology can be particularly advantageous to older adults, patients with long-term illnesses, athletes, and people who wish to lead a healthy life. Wearable technology prompts people to remain physically active through notifications for physical activities, hydration, taking their prescribed medications, and sleeping adequately. In times of emergencies, some sophisticated wearables can detect any unusual behavior such as falling down, abnormal heartbeat, and even vital sign abnormalities. These innovations provide greater convenience, fewer hospital visits, and help promote preventive medicine.Track heart rate, activity, sleep patterns.



Figure 5. Data flow process in IoT-enabled healthcare system.

2.3 Smart Hospitals

The concept of smart hospitals involves the implementation of technological advances within the framework of a contemporary healthcare facility. Specifically, IoT, AI, cloud computing, robotic solutions, and data analytics can be used to enhance patient experience and improve hospital administration and performance. A smart hospital has various interconnected devices like sensors, smart beds, patient monitoring systems, RFID devices, and robots generating data through secure connections. The medical professionals can control patients' heart rate, blood pressure, body temperature, and oxygen levels in real-time, thus making their treatment more effective. Moreover, smart hospitals apply digital solutions to schedule appointments, maintain health records electronically, provide telemedicine services, trace medication usage, and manage emergencies. Finally, the IoT technologies allow identifying and managing the location and functioning of the hospital's equipment, contributing to its better utilization. Besides, the smart hospital has automated lighting, air-conditioning, and energy management systems.

The use of AI can help in diagnosing diseases, analyzing images, and predicting patients' risk factors. The benefits of a smart hospital include reduction in paper work, fewer human errors, and enhanced communication among health professionals. The advantages for patients include reduced wait times, increased safety during treatment, personalization of healthcare services, and improved patient experience. In case of an emergency or pandemic, smart hospitals can facilitate tele-consultation and effective patient management.

It includes smart beds, asset tracking, and automation.



Figure 6. Smart Hospitals

2.4 Telemedicine

Telemedicine involves the provision of healthcare services through electronic technology. In this case, physicians and specialists offer remote consultation services, diagnosis, monitoring, and treatment of various illnesses without seeing the patients physically. Telemedicine involves the utilization of technologies such as IoT, video conferencing, mobile applications, and digital communication technology to facilitate the delivery of healthcare services. Patients can share their symptoms with physicians and get advice on how to proceed, prescriptions for medicines, and further consultation. Physicians are able to access information about the patient including images and live data gathered from wearable technologies like heart rate monitors, blood glucose levels, blood pressure, and pulse oximeters.

Telemedicine proves extremely helpful for those residing in the rural or remote parts of the country because it cuts down the time of travelling to the hospital and makes healthcare less costly and convenient. In cases of emergency situations like pandemics, telemedicine proves extremely important due to its capability to ensure consultation without putting both the doctor and the patient at the risk of infection.

Telemedicine also helps in providing consistent treatment for the elderly patients as well as those suffering from any sort of chronic illness by maintaining constant contact with them. Telemedicine is a blend of healthcare and communication technology that helps in better provision of medical care.

Enables remote consultation and diagnosis.



Figure 7. Wearable IoT system for continuous patient monitoring

3. BENEFITS OF IOT IN HEALTHCARE

Innovation in technology has seen revolutionary changes in the healthcare industry due to the connection between medical instruments, sensors, software programs, and healthcare workers via the internet. The IoT has made possible effective data gathering, communication, and decision making, resulting in better healthcare service delivery. One of the major advantages of IoT in healthcare is remote monitoring of patients. Medical devices like heart rate monitors, blood pressure monitors, glucose level detectors, pulse oximeters, and even wearable fitness bands are able to gather patient data and transmit it to doctors and hospitals where the data is monitored.

Early detection of illnesses and preventative care is yet another significant advantage of IoT. The Internet of Things can detect any early symptoms of an illness. If one suffers from heart rate disturbances, lack of oxygen in his body, or even a sudden increase in blood pressure, IoT will detect them in time and help to cure these disorders effectively.

IoT enables remote health monitoring for elderly patients, chronically ill people, and even people who had recently undergone surgery. Such people can have medical surveillance at home instead of going to the hospital.

Hospital operations will be managed more efficiently due to IoT. With the help of IoT technology, smart hospitals manage their patients better, schedule their appointments, manage their medicines, and track all their assets and equipment. Wheelchairs, ventilators, infusion pumps, and other medical appliances may be found faster with the help of IoT and RFID tags. Lighting and temperature can also be controlled automatically.

Telemedicine is yet another key advantage provided by the IoT framework. Telemedicine involves consultations with patients via video conferencing and access to data generated by devices connected to the IoT network. Such an approach would prove highly advantageous for remote regions where specialist medical facilities are few in number. In doing so, both time and money saved.

Moreover, IoT technology also contributes significantly to enhancing data accuracy and minimizing the scope of human error. The automatic generation of patient-related data makes human error almost nil in nature. The availability of real-time and accurate data would enable doctors to take more effective decisions. Further, IoT systems could even send reminders for medicines, appointments, and physical activities.

In case of pandemics or emergencies, IoT has been immensely useful due to its ability to offer contactless monitoring and remote healthcare services. Even though data privacy and security is still an issue, appropriate measures can resolve these issues. IoT has made healthcare safer, effective, efficient, accessible, and cost-effective.

Some another benefit of IoT in Healthcare

- Real-time monitoring
- Improved patient care
- Cost reduction
- Enhanced efficiency
- Data-driven decisions

4. CHALLENGES OF IOT

The IoT provides numerous benefits in the healthcare sector and other industries; however, some significant problems exist, which might hinder the deployment of the IoT successfully. The problem of data security and privacy is one of the biggest concerns. The IoT devices gather and send personal information about patients and their health status constantly. If adequate security policies are not implemented, then hackers can get access to confidential data. Hence, it is vital to implement encryption techniques and other security strategies.

Interoperability is another major problem related to the IoT. Sometimes, various devices developed by diverse vendors utilize different standards and protocols for communication. As a result, devices cannot interact easily with one another. To address this issue, standardization needs to be achieved.

Issues of reliability and connectivity also play a vital role. IoT devices are dependent on consistent internet and power supply. In case of inadequate internet network coverage in rural or distant areas, there could be an issue regarding interruption in data transfer. Failure of medical IoT devices would affect patient monitoring.

The high cost involved in implementation and maintenance is another potential drawback. Sensors, gateways, cloud computing platform, software solutions, and networking security require considerable financial expenditure. Smaller hospitals or health care facilities may find it hard to implement such technologies due to high costs. Additionally, maintenance and updates would also add to expenses.

Managing data is yet another issue due to the enormous amount of data generated by IoT devices per second. For this purpose, strong servers, cloud computing, and sophisticated analytical software are needed. Otherwise, valuable data might not be retrieved.

Short battery life and poor durability may cause difficulties, particularly when it comes to wearable devices or remote sensors. Continuous charging and frequent replacements decrease the comfort level. Lastly, insufficient technical expertise and user knowledge could impede adoption of the technology. It is necessary to train physicians, nurses, and patients on how to utilize IoT technology efficiently.

Another benefits are:

- Data security
- Interoperability
- High cost
- Data overload
- Regulatory issues



Figure 8. IoT-enabled smart hospital ecosystem

5. ETHICAL CONSIDERATIONS

It is important to consider ethics in the application of IoT technology, especially in the medical sector as there is a risk of misuse and abuse of sensitive information related to patient confidentiality. The first ethical issue is that of privacy since IoT devices like wearable, monitors, and medical devices gather personal information such as heartbeat, blood pressure, location, and even activities of a day. Individuals must be made aware of what data they are providing, its uses, and who might access it to ensure confidentiality is observed.

The other ethical issue that must be considered is informed consent. There is need for users to give voluntary consent after being made aware of benefits and possible risks associated with the collection of data.

Patients in medical facilities must know what happens to their data as regards sharing it with physicians, medical institutions, insurance companies, or researchers.

Security is another responsibility when using these technologies. The failure of any health care institution to guard the system against malicious attacks may result in the patients' discrimination or injury. This means that adequate security measures such as encryption, good password protection, and updating software should be put in place. The accuracy of the data is also a challenge since inaccurate data collected through these devices could result in an incorrect diagnosis or treatment decisions.

Access to technology is also a responsibility that needs to be taken seriously. The use of such advanced IoT technologies in healthcare may be too costly and unavailable in remote areas, especially rural areas or developing countries. It should ensure equality of access to the technology regardless of people's financial capability. There should also be considerations concerning reliance on artificial intelligence or automation.

Accountability should also be established where things go wrong. Responsibility must be determined regarding the mistakes made in the usage of the technology by individuals or organizations involved.

Other benefits:

- Privacy
- Data ownership
- Accountability

6. FUTURE TRENDS

The future of the IoT in healthcare promises to be bright due to a number of technological developments that can lead to additional innovations in the field. One of the main future directions involves the utilization of Artificial Intelligence (AI) and Machine Learning (ML) in conjunction with IoT-enabled equipment. These systems would have the ability to process patient data on a continuous basis and diagnose illnesses early while offering customized treatment options. An AI-driven medical aid would assist physicians in making quick and efficient decisions.

An additional trend would be the development of wearables and implantables in medical technology. The next generation of wearables would be compact, comfortable, and accurate, constantly tracking vital signs such as heartbeat, blood glucose level, blood pressure, sleep patterns, and others. Implantable sensors can monitor internal body functions and alert doctors about abnormalities without their intervention. These tools would contribute to preventive medicine and chronic illness management.

Another factor would be the spread of 5G technology and efficient communication systems. The fast connection, minimum delays, and dependable connection would allow surgeons to operate remotely and communicate with patients effectively. Smart ambulances in connection with hospitals could send patient information to doctors in advance to allow for pre-planning of emergency interventions.

Cloud computing and edge computing will increasingly be utilized to process massive amounts of healthcare information efficiently. The edge computing system can perform data processing locally within the device, which eliminates any delay and increases the response speed. The blockchain technique might be implemented to ensure security and transparency when sharing the patient's personal health records.

The future smart hospitals will have robotic assistance, automated medication delivery systems, smart beds, and other intelligent management systems. Future personalized medicine will utilize real-time data from IoT devices to achieve better medical results. Even though security and regulatory compliance will remain key concerns, future trends in IoT in healthcare will lead to smarter, more efficient, secure, and patient-centric healthcare around the globe.

Some other advantages are:

- AI integration
- 5G technology
- Blockchain
- Smart implants
- Personalized medicine

Figure 9. Security architecture for IoT healthcare systems

CONCLUSION

The Internet of Things (IoT) has emerged as one of the greatest technologies that have revolutionized the healthcare industry by linking medical devices, sensors, software applications, and healthcare professionals through the internet. The IoT technology helps in enabling real-time monitoring, faster communication, information gathering, and intelligent decision-making, thereby improving the level of care for patients. IoT has brought about innovation in the health care industry through telemedicine platforms, remote monitoring solutions, smart hospitals, wearable devices, among others.

The Internet of Things (IoT) is critical in transforming the medical sector from the current approach of reactive care to a proactive and preventive one. IoT devices enable continuous monitoring of physiological parameters such as heartbeat, blood pressure, oxygen concentration, and glucose levels to facilitate early identification of any irregularities, leading to timely medical attention.

This strategy minimizes hospital visits, cuts down medical expenses, and enhances patient outcomes. Additionally, patients have the luxury of being under constant medical surveillance and consultation from their residential premises.

Although IoT provides several benefits, there are various difficulties that it encounters, including issues such as data security, cybersecurity threats, lack of interoperability, exorbitant setup costs, and regulation. Ethics such as obtaining consent, equality, and accountability are significant concerns that should not be overlooked.

Looking into the future, the prospects for IoT in medical sciences seem extremely bright. The use of AI, machine learning, 5G technology, cloud computing, and smart analytics is bound to lead to even greater advancements in the sphere of healthcare. In addition, hospitals and clinics are likely to be transformed into truly smart facilities with individualized and precise approaches to treatment. In conclusion, IoT is transforming healthcare through increased efficiency, safety, and accessibility.

REFERENCES

Alnajim, R., &Alkhalifah, A. (2025). Exploring factors affecting the adoption of IoT in healthcare: A systematic literature review. Healthcare, 13(23), 3157. https://www.mdpi.com/2227-9032/13/23/3157

Al-rawashdeh, M., Keikhosrokiani, P., Belaton, B., Alawida, M., &Zwiri, A. (2022). IoT adoption and application for smart healthcare: A systematic review. Sensors, 22(14), 5377. https://www.mdpi.com/1424-8220/22/14/5377

Bacancy Technology. (2023). IoT in healthcare: Benefits, challenges & future trends. https://www.bacancytechnology.com

Digi International. (2023). IoT in healthcare applications. https://www.digi.com

HData Systems. (2022). IoT in healthcare: Benefits and future. https://www.hdatasystems.com

MDPI. (2024). IoT applications and challenges in global healthcare systems: A comprehensive review. https://www.mdpi.com

N-iX. (2022). IoT healthcare benefits and challenges. https://www.n-ix.com

ValueCoders. (2023). IoT in healthcare benefits and challenges. https://www.valuecoders.com

Zhang, Y., Yang, G., & Wang, L. (2021). A systematic review of IoT in healthcare: Applications, techniques, and trends. Journal of Network and Computer Applications, 192, 103164.

Zhou, X., Li, Q., & Chen, W. (2024). A review of IoT applications in healthcare. Neurocomputing, 565, 127017.



ISBN: 978-625-90649-0-1